

2A-orbifold construction and the baby-monster vertex operator superalgebra

Hiroshi Yamauchi

Graduate School of Mathematics,

University of Tsukuba, Ibaraki 305-8571, Japan

e-mail: hirocci@math.tsukuba.ac.jp

Abstract

In this article we prove that the full automorphism group of the baby-monster vertex operator superalgebra constructed by Höhn is isomorphic to $2 \times \mathbb{B}$, where \mathbb{B} is the baby-monster sporadic finite simple group, and determine irreducible modules for the baby-monster vertex operator algebra. Our result has many corollaries. In particular, we can prove that the \mathbb{Z}_2 -orbifold construction with respect to a 2A-involution of the Monster applied to the moonshine vertex operator algebra V^\natural yields V^\natural itself again.

1 Introduction

The famous moonshine vertex operator algebra V^\natural constructed by Frenkel-Lepowsky-Murman [FLM] is the first example of the \mathbb{Z}_2 -orbifold construction of a holomorphic vertex operator algebra (VOA). Let us explain a \mathbb{Z}_2 -orbifold construction briefly. Let V be a holomorphic vertex operator algebra and σ an involutive automorphism on V . Then the fixed point subalgebra $V^{(\sigma)}$ is a simple vertex operator algebra. It is shown in [DLM1] that there is a unique irreducible σ -twisted V -module M and we have a decomposition $M = M^0 \oplus M^1$ into a direct sum of irreducible $V^{(\sigma)}$ -modules such that M^0 has an integral top weight. Then a \mathbb{Z}_2 -orbifold construction with respect to $\sigma \in \text{Aut}(V)$ is to construct a \mathbb{Z}_2 -graded extension $W = V^{(\sigma)} \oplus M^0$ of the fixed point subalgebra $V^{(\sigma)}$ which is expected to be a holomorphic vertex operator algebra.

In FLM's construction, we take V to be the lattice vertex operator algebra V_Λ associated to the Leech lattice Λ and the involution σ is a natural lifting $\theta \in \text{Aut}(V_\Lambda)$ of the (-1) -isometry on Λ . Denote by $V_\Lambda = V_\Lambda^+ \oplus V_\Lambda^-$ the eigenspace decomposition such that θ acts on V_Λ^\pm as ± 1 . Let V_Λ^T be the unique irreducible θ -twisted V_Λ -module. Then there is a decomposition $V_\Lambda^T = (V_\Lambda^T)^+ \oplus (V_\Lambda^T)^-$ such that the top weight of $(V_\Lambda^T)^+$ is integral. Then

the moonshine vertex operator algebra is defined by $V^\natural := V_\Lambda^+ \oplus (V_\Lambda^T)^+$ and it is proved in [FLM] that V^\natural forms a \mathbb{Z}_2 -graded extension of V_Λ^+ . It is also proved in [FLM] that the full automorphism group of the moonshine vertex operator algebra is the Monster sporadic finite simple group \mathbb{M} by using Griess' result [G].

In the Monster, there are two conjugacy classes of involutions, the 2A-conjugacy class and the 2B-conjugacy class (cf. [ATLAS]). One can explicitly see the action of a 2B-involution on V^\natural by FLM's construction. But it is not clear to see the action of a 2A-involution on V^\natural before Miyamoto. In [M1], Miyamoto opened a way to study the action of 2A-involutions of the Monster on the moonshine VOA by using a sub VOA isomorphic to the unitary Virasoro VOA $L(1/2, 0)$. Let us recall the definition of Miyamoto involutions. Let V be a simple VOA and $e \in V_2$ be a vector such that e generates a sub VOA isomorphic to $L(1/2, 0)$. Such a vector e is called conformal vector with central charge $1/2$. Since V as a $\text{Vir}(e) \simeq L(1/2, 0)$ -module is completely reducible, we have a decomposition

$$V = V_e(0) \oplus V_e(1/2) \oplus V_e(1/16),$$

where $V_e(h)$ denotes a sum of all irreducible $\text{Vir}(e)$ -submodules isomorphic to $L(1/2, h)$, $h = 0, 1/2, 1/16$. Then one can define a linear isomorphism τ_e on V by

$$\tau_e := 1 \quad \text{on } V_e(0) \oplus V_e(1/2), \quad -1 \quad \text{on } V_e(1/16).$$

Then it is proved in [M1] that τ_e defines an involution of a VOA V if $V_e(1/16) \neq 0$. This involution is often called the (first) Miyamoto involution. If $V_e(1/16) = 0$, then one can define another automorphism on V by

$$\sigma_e := 1 \quad \text{on } V_e(0), \quad -1 \quad \text{on } V_e(1/2).$$

This involution is also called the (second) Miyamoto involution. It is shown in [C] and [M1] that in the moonshine VOA every Miyamoto involution τ_e defines a 2A-involution of the Monster and the correspondence between conformal vectors and 2A-involutions is one-to-one. Therefore, in the study of 2A-involutions, it is very important to study conformal vectors with central charge $1/2$. Along this idea, C.H. Lam, H. Yamada and the author obtained an interesting achievement on 2A-involutions of the Monster in [LYY].

The main purpose of this paper is to study the \mathbb{Z}_2 -orbifold construction of V^\natural with respect to the Miyamoto involution and to prove that the 2A-orbifold construction applied to V^\natural yields V^\natural itself again. Since a 2A-involution of the Monster is uniquely determined by a conformal vector e of V^\natural with central charge $1/2$, we first study the commutant subalgebra of $\text{Vir}(e)$. For a simple VOA V and a conformal vector e of V with central charge $1/2$, set the space of highest weight vectors by $T_e(h) := \{v \in V \mid e_{(1)}v\}$ for $h = 0, 1/2, 1/16$. Then we have decompositions $V_e(h) = L(1/2, h) \otimes T_e(h)$ and the

commutant subalgebra $T_e(0)$ acts on $T_e(h)$ for $h = 0, 1/2, 1/16$. Since $L(1/2, 0)$ has a \mathbb{Z}_2 -graded extension $L(1/2, 0) \oplus L(1/2, 1/2)$, we can introduce a vertex operator superalgebra (SVOA) structure on $T_e(0) \oplus T_e(1/2)$ (Theorem 3.6) and its \mathbb{Z}_2 -twisted module structure on $T_e(1/16)$ (Theorem 3.8). It is easy to see that the one point stabilizer $C_{\text{Aut}(V)}(e)$ of a conformal vector e naturally acts on the space of highest weight vectors $T_e(h)$. If we take $V = V^\natural$, then $C_{\text{Aut}(V^\natural)}(e)$ is isomorphic to the 2-fold central extension $\langle \tau_e \rangle \cdot \mathbb{B}$ of the baby-monster sporadic finite simple group \mathbb{B} . Therefore, the SVOA $T_e^\natural(0) \oplus T_e^\natural(1/2)$, where we have set $V_e^\natural(h) = L(1/2, h) \otimes T_e^\natural(h)$ for $h = 0, 1/2, 1/16$, affords a natural action of \mathbb{B} . Motivated by this fact, Höhn first studied this SVOA in [Hö1] and he called it the baby-monster SVOA. Following him, we write $VB^0 := T_e^\natural(0)$, $VB^1 := T_e^\natural(1/2)$ and $VB := T_e^\natural(0) \oplus T_e^\natural(1/2)$. It is proved in [Hö2] that the full automorphism group of the even part VB^0 of VB is exactly isomorphic to the baby-monster \mathbb{B} . In this paper, we give a quite different proof of $\text{Aut}(VB^0) \simeq \mathbb{B}$ based on a theory of simple current extensions.

In my recent work [Y1] [Y2], a theory of simple current extensions of vertex operator algebras was developed and many useful results were obtained. Using this theory, we determine the automorphism group of the commutant subalgebra $T_e(0)$ as follows:

Theorem 1. *Let V be a holomorphic VOA and $e \in V$ a conformal vector with central charge $1/2$. Suppose the following:*

- (a) $V_e(h) \neq 0$ for $h = 0, 1/2, 1/16$,
- (b) $V_e(0)$ and $T_e(0)$ are rational C_2 -cofinite VOAs of CFT-type,
- (c) $V_e(1/16)$ is a simple current $V^{\langle \tau_e \rangle}$ -module,
- (d) $T_e(1/2)$ is a simple current $T_e(0)$ -module,
- (e) $C_{\text{Aut}(V)}(e)/\langle \tau_e \rangle$ is a simple group or an odd group.

Then

- (i) $\text{Aut}(T_e(0)) = C_{\text{Aut}(V)}(e)/\langle \tau_e \rangle$.
- (ii) The irreducible $T_e(0)$ -modules are given by $T_e(0)$, $T_e(1/2)$ and $T_e(1/16)$.
- (iii) The τ_e -orbifold construction applied to V yields V itself again.

The assumptions (c) and (d) in the theorem above seem to be rather restrictive. However, we prove that all the assumptions above hold if V is the moonshine VOA. We also present a refinement of Miyamoto's reconstruction of the moonshine VOA [M5]. Our refinement enable us to prove not only that the baby-monster SVOA VB satisfies all the assumptions above but also that we can construct the baby-monster SVOA VB without reference to V^\natural . The main theorem of this paper is

Theorem 2. *Let $VB = VB^0 \oplus VB^1$ the simple SVOA obtained from V^\natural .*

- (1) $\text{Aut}(VB^0) = \mathbb{B}$ and $\text{Aut}(VB) = 2 \times \mathbb{B}$.
- (2) There are exactly three irreducible VB^0 -modules, VB^0 , VB^1 and $VB_T := T_e^\natural(1/16)$.

(3) The fusion rules for VB^0 -modules are as follows:

$$VB^1 \times VB^1 = VB^0, \quad VB^1 \times VB_T = VB_T, \quad VB_T \times VB_T = VB^0 + VB^1.$$

This theorem has many corollaries:

Corollary 1. *The irreducible $2A$ -twisted V^\natural -module has a shape*

$$L(1/2, 1/2) \otimes VB^0 \oplus L(1/2, 0) \otimes VB^1 \oplus L(1/2, 1/16) \otimes VB_T.$$

Corollary 2. *For any conformal vector $e \in V^\natural$ with central charge $1/2$, there is no $\rho \in \text{Aut}(V^\natural)$ such that $\rho(V_e^\natural(h)) = V_e^\natural(h)$ for $h = 0, 1/2, 1/16$ and $\rho|_{(V^\natural)^{\langle \tau_e \rangle}} = \sigma_e$.*

Corollary 3. *The $2A$ -orbifold construction applied to the moonshine VOA V^\natural yields V^\natural itself again.*

At the end of this paper, we give characters of VB^0 -modules and their modular transformation laws. Surprisingly, we find that the fusion algebra and the modular transformation laws for the baby-monster VOA is canonically isomorphic to those of the Ising model $L(1/2, 0)$.

2 Simple current extension

Let V be a simple vertex operator algebra (VOA). We recall a definition of a fusion product of V -modules.

Definition 2.1. Let M^1, M^2 be V -modules. A *fusion product* for the ordered pair (M^1, M^2) is a pair $(M^1 \boxtimes_V M^2, F(\cdot, z))$ consisting of a V -module $M^1 \boxtimes_V M^2$ and a V -intertwining operator $F(\cdot, z)$ of type $M^1 \times M^2 \rightarrow M^1 \boxtimes_V M^2$ satisfying the following universal property: For any V -module W and any V -intertwining operator $I(\cdot, z)$ of type $M^1 \times M^2 \rightarrow W$, there exists a unique V -homomorphism ψ from $M^1 \boxtimes_V M^2$ to W such that $I(\cdot, z) = \psi F(\cdot, z)$. We usually denote the pair $(M^1 \boxtimes_V M^2, F(\cdot, z))$ simply by $M^1 \boxtimes_V M^2$.

A theory of fusion products has been greatly developed by Huang-Lepowsky [HL1]-[HL4] and Huang [H1]-[H4] (see also [DLM2] [Li3]), and it is proved that if V is rational then a fusion product of any two V -modules always exists (cf. [HL3] [HL4] [Li3]) and if V is also C_2 -cofinite and of CFT-type then the fusion product satisfies the associativity (cf. [H1] [H4] [DLM2]). Therefore, the theory of fusion products is a powerful tool to study a rational C_2 -cofinite vertex operator algebra of CFT-type. Among modules for such a vertex operator algebra, simple current modules have a special importance.

Definition 2.2. An irreducible V -module U is called a *simple current* if it satisfies: For any irreducible V -module W , the fusion product $U \boxtimes_V W$ is also irreducible.

In this paper we mainly consider the following extensions of vertex operator algebras.

Definition 2.3. Let V^0 be a simple rational VOA and D a finite abelian group. Let $\{V^\alpha \mid \alpha \in D\}$ be a set of inequivalent irreducible V^0 -modules. A D -graded extension V_D of V^0 is a simple VOA $V_D = \bigoplus_{\alpha \in D} V^\alpha$ which extends the VOA structure of V^0 with the grading structure $Y_{V_D}(x^\alpha, z)x^\beta \in V^{\alpha+\beta}((z))$ for any $x^\alpha \in V^\alpha$, $x^\beta \in V^\beta$. A D -graded extension V_D is called a D -graded simple current extension of V^0 if all V^α , $\alpha \in D$, are simple current V^0 -modules. In the case of $D = \mathbb{Z}_2 = \{0, 1\}$ and $V_D = V^0 \oplus V^1$ is a simple vertex operator superalgebra with even part V^0 and odd part V^1 , we call V_D a *simple current super-extension* of V^0 if V^1 is a simple current V^0 -module.

Remark 2.4. Let D^* be the dual group of D . By definition, there is a natural action of D^* on V_D defined as $\chi|_{V^\alpha} = \chi(\alpha) \cdot \text{id}_{V^\alpha}$ for $\alpha \in D$ and $\chi \in D^*$.

We have the following uniqueness property of a simple current extension.

Lemma 2.5. ([DM2]) (i) Let $V_D = \bigoplus_{\alpha \in D} V^\alpha$ be a D -graded extension of V^0 . If the space of V^0 -intertwining operators of type $V^\alpha \times V^\beta \rightarrow V^{\alpha+\beta}$ is one-dimensional for all $\alpha, \beta \in D$, then the VOA structure on V_D is unique over \mathbb{C} . In particular, if V_D is a D -graded simple current extension of V^0 , then its VOA structure is unique over \mathbb{C} .
(ii) The SVOA structure on a simple current super-extension is unique over \mathbb{C} .

In general, it is a difficult problem to determine whether a given module is a simple current or not. However, the following lemma provides us a simple criterion.

Lemma 2.6. ([Y2]) Let V be a simple rational C_2 -cofinite VOA of CFT-type and U a V -module. If there is a V -module W such that the fusion rule $U \boxtimes_V W = V$ holds in the fusion algebra for V , then U (and also W) is a simple current V -module.

Proof: By the assumption, we can use the results in [H4] so that the fusion algebra for V is a commutative associative algebra over \mathbb{N} with the unit element V . Let X be an irreducible V -module. Then $U \boxtimes_V X$ is also a V -module and is a direct sum of irreducible V -modules. Let $U \boxtimes_V X = \bigoplus_{i \in I} T^i$ be a decomposition into a direct sum of irreducible V -modules. First, we show that $W \boxtimes_V T^i \neq 0$ for all $i \in I$. Assume that there is an $i_0 \in I$ such that $W \boxtimes_V T^{i_0} = 0$. Then by multiplying U in the fusion algebra we obtain $0 = U \boxtimes_V (W \boxtimes_V T^{i_0}) = (U \boxtimes_V W) \boxtimes_V T^{i_0} = V \boxtimes_V T^{i_0} = T^{i_0}$, a contradiction. Therefore, $W \boxtimes_V T^i \neq 0$ for all $i \in I$. Then by multiplying W to $U \boxtimes_V X$ we get $X = V \boxtimes_V X = (W \boxtimes_V U) \boxtimes_V X = W \boxtimes_V (U \boxtimes_V X) = W \boxtimes_V \bigoplus_{i \in I} T^i = \bigoplus_{i \in I} W \boxtimes_V T^i$.

Therefore, the cardinality of the index set I is 1 and hence $U \boxtimes_V X$ is an irreducible V -module. Thus U is a simple current V -module. \blacksquare

The representation theory of simple current extensions is studied in many papers (cf. [DLM3] [L1] [SY] [Y1] [Y2]) and it is shown in [L1] [Y1] [Y2] that every simple current extension of a simple rational C_2 -cofinite VOA of CFT-type is also rational and C_2 -cofinite. We review some results from [Y1] and [Y2] which we will need later.

Let $V_D = \bigoplus_{\alpha \in D} V^\alpha$ be a D -graded simple current extension of a simple rational C_2 -cofinite CFT-type VOA V^0 . Since the fusion algebra for V^0 is associative, we can adopt the following definition.

Definition 2.7. Let W be an irreducible V^0 -module W . A subset $D_W := \{\alpha \in D \mid V^\alpha \boxtimes_V W \simeq W\}$ forms a subgroup of D . We call D_W the *stabilizer* of W .

Lemma 2.8. ([SY]) Let M be a V_D -module and W an irreducible V^0 -submodule of M . Then $V^\alpha \cdot W$ is also a non-trivial irreducible V^0 -submodule of M , where $V^\alpha \cdot W$ denotes a linear space spanned by elements $a_{(n)}w = \text{Res}_z z^n Y_M(a, z)w$ with $a \in V^\alpha$, $w \in W$ and $n \in \mathbb{Z}$. The stabilizer D_W is determined independently of a choice of an irreducible V^0 -submodule W if M is an irreducible V_D -module.

By the lemma above, we introduce the following notion.

Definition 2.9. An irreducible V_D -module M is called *D -stable* if $D_W = 0$ for some irreducible V^0 -submodule W of M .

Among V_D -modules, D -stable modules enjoy nice properties.

Proposition 2.10. ([SY] [Y1]) Let M be an irreducible D -stable V_D -module. Then the V_D -module structure on a V^0 -module M is unique over \mathbb{C} .

Theorem 2.11. ([Y1], Induced modules) Let W be an irreducible V^0 -module. Then there exists a unique $\chi \in D^*$ such that W is contained in an irreducible χ -twisted V_D -module. If $D_W = 0$, then it is given by the induced module

$$\text{Ind}_{V^0}^{V_D} W := \bigoplus_{\alpha \in D} V^\alpha \boxtimes_{V^0} W.$$

Moreover, any irreducible χ -twisted V_D -module containing W as a V^0 -submodule is isomorphic to $\text{Ind}_{V^0}^{V_D} W$ above.

For a later purpose, we give a detailed description of the theorem above in the case of simple current super-extensions.

Theorem 2.12. ([Y2]) Let $V = V^0 \oplus V^1$ be a simple current super-extension of a simple rational C_2 -cofinite VOA V^0 of CFT-type. For an irreducible V^0 -module W ,

(i) If $V^1 \boxtimes_{V^0} W \not\simeq W$, then W is uniquely lifted to be an irreducible untwisted or \mathbb{Z}_2 -twisted V -module $W \oplus (V^1 \boxtimes_{V^0} W)$.

(ii) If $V^1 \boxtimes_{V^0} W \simeq W$, then there exist exactly two inequivalent irreducible \mathbb{Z}_2 -twisted V -module structure on W . If we write one of them by W^+ , then the other one is given as the \mathbb{Z}_2 -conjugate V -module of W^+ .

Theorem 2.13. ([SY] [Y1], Lifting property of intertwining operators) Let $M^i, i = 1, 2, 3$, be irreducible D -stable V_D -modules and let W^i be irreducible V^0 -submodules of M^i for $i = 1, 2, 3$. Then for a V^0 -intertwining operator $I(\cdot, z)$ of type $W^1 \times W^2 \rightarrow W^3$ there is a V_D -intertwining operator $\tilde{I}(\cdot, z)$ of type $M^1 \times M^2 \rightarrow M^3$ such that $\tilde{I}(\cdot, z)|_{W^1 \otimes W^2} = I(\cdot, z)$. Therefore, there is a natural linear isomorphism

$$\begin{pmatrix} M^3 \\ M^1 \ M^2 \end{pmatrix}_{V_D} \simeq \bigoplus_{\alpha \in D} \begin{pmatrix} V^\alpha \boxtimes_{V^0} W^3 \\ W^1 \quad W^2 \end{pmatrix}_{V^0},$$

where $\begin{pmatrix} M^3 \\ M^1 \ M^2 \end{pmatrix}_{V_D}$ denotes the space of V_D -intertwining operators of type $M^1 \times M^2 \rightarrow M^3$.

Corollary 2.14. Let E be a subgroup of D and fix a coset decomposition $D = \sqcup_{i=1}^r (\alpha_i + E)$ of D . Then $V_E := \bigoplus_{\alpha \in E} V^\alpha$ is an E -graded simple current extension of V^0 and $V_{E+\alpha_i} := \bigoplus_{\beta \in E} V^{\alpha_i+\beta}$ are simple current V_E -modules. Therefore, we can view $V_D = \bigoplus_{i=1}^r V_{E+\alpha_i}$ as a D/E -graded simple current extension of V_E .

Let us consider automorphisms on V_D . Let $\sigma \in \text{Aut}(V^0)$ and $(X, Y_X(\cdot, z))$ a V^0 -module. We can define the σ -conjugate module X^σ as follows. As a vector space, we set $X^\sigma = X$ and the vertex operator map on X^σ is defined by $Y_{X^\sigma}(a, z) := Y_X(\sigma a, z)$ for $a \in V^0$. It is clear that X^σ is irreducible if and only if X is an irreducible V^0 -module. An irreducible V^0 -module X is called σ -stable if its σ -conjugate X^σ is isomorphic to X as a V^0 -module.

Remark 2.15. The definition of D -stable V_D -modules comes from the fact that if M is an irreducible D -stable V_D -module then its conjugate M^χ is isomorphic to M as a V_D -module for any $\chi \in D^*$.

We can construct the σ -conjugate $(V_D)^\sigma$ of V_D as follows. Let $Y_{V_D}(\cdot, z)$ be the vertex operator map on V_D . By definition, there are canonical linear isomorphisms $\psi_\alpha : V^\alpha \rightarrow (V^\alpha)^\sigma, \alpha \in D$, such that

$$Y_{(V^\alpha)^\sigma}(x^0, z)\psi_\alpha = \psi_\alpha Y_{V^\alpha}(\sigma x^0, z)$$

for all $x^0 \in V^0$. Then we define the vertex operator map $Y_{V_D}^\sigma(\cdot, z)$ on $(V_D)^\sigma = \bigoplus_{\alpha \in D} (V^\alpha)^\sigma$ by

$$Y_{V_D}^\sigma(\psi_\alpha x^\alpha, z)\psi_\beta := \psi_{\alpha+\beta} Y_{V_D}(x^\alpha, z)x^\beta$$

for $x^\alpha \in V^\alpha$, $x^\beta \in V^\beta$. Then one can easily verify that $((V_D)^\sigma, Y_{V_D}^\sigma(\cdot, z))$ also forms a D -graded simple current extension of V^0 . The following lifting property is established in [Sh] by using the uniqueness of VOA structure on V_D .

Theorem 2.16. (*[Sh], Lifting property of automorphisms*) *Let $\sigma \in \text{Aut}(V^0)$ such that $(V_D)^\sigma \simeq V_D$ as a V^0 -module. Then there is a lifting $\tilde{\sigma} \in \text{Aut}(V_D)$ such that $\tilde{\sigma}V^0 = V^0$ and $\tilde{\sigma}|_{V^0} = \sigma$. The lifting $\tilde{\sigma}$ is unique up to multiples of elements in $D^* \subset \text{Aut}(V_D)$. This assertion still holds if $D = \mathbb{Z}_2$ and $V_D = V^0 \oplus V^1$ is a simple current super-extension of V^0 .*

As we have seen above, simple current extensions have many good properties. At last of this section, we present the following extension property of simple current extensions.

Theorem 2.17. (*[Y2], Extension property of simple current extensions*) *Let $V^{(0,0)}$ be a simple rational C_2 -cofinite VOA of CFT-type, and let D_1, D_2 be finite abelian groups. Assume that we have a set of inequivalent irreducible simple current $V^{(0,0)}$ -modules $\{V^{(\alpha,\beta)} \mid (\alpha, \beta) \in D_1 \oplus D_2\}$ with $D_1 \oplus D_2$ -graded fusion rules $V^{(\alpha_1, \beta_1)} \boxtimes_{V^{(0,0)}} V^{(\alpha_2, \beta_2)} = V^{(\alpha_1 + \alpha_2, \beta_1 + \beta_2)}$ for any $(\alpha_1, \beta_1), (\alpha_2, \beta_2) \in D_1 \oplus D_2$.*

(i) *Further assume that all $V^{(\alpha, \beta)}$, $(\alpha, \beta) \in D_1 \oplus D_2$, have integral top weights and we have D_1 - and D_2 -graded simple current extensions $V_{D_1} = \bigoplus_{\alpha \in D_1} V^{(\alpha, 0)}$ and $V_{D_2} = \bigoplus_{\beta \in D_2} V^{(0, \beta)}$. Then $V_{D_1 \oplus D_2} := \bigoplus_{(\alpha, \beta) \in D_1 \oplus D_2} V^{(\alpha, \beta)}$ possesses a unique structure of a simple vertex operator algebra as a $D_1 \oplus D_2$ -graded simple current extension of $V^{(0,0)}$.*

(ii) *In the case of $D_2 = \mathbb{Z}_2 = \{0, 1\}$, further assume that all $V^{(\alpha, 0)}$, $\alpha \in D_1$, have integral top weight, all $V^{(\alpha, 1)}$, $\alpha \in D_1$, have half-integral top weight, $V_{D_1} = \bigoplus_{\alpha \in D_1} V^{(\alpha, 0)}$ is a D_1 -graded simple current extension of $V^{(0,0)}$, and $V_{D_2} = V^{(0,0)} \oplus V^{(0,1)}$ is a simple current super-extension of $V^{(0,0)}$. Then $V_{D_1 \oplus D_2} = \bigoplus_{(\alpha, \beta) \in D_1 \oplus D_2} V^{(\alpha, \beta)}$ has a unique structure of a simple vertex operator superalgebra with even part $\bigoplus_{\alpha \in D_1} V^{(\alpha, 0)}$ and odd part $\bigoplus_{\beta \in D_1} V^{(\beta, 1)}$ as a simple current super-extension of V_{D_1} .*

3 Miyamoto involution and its centralizer

Let us denote by $L(c, h)$ the irreducible highest weight module for the Virasoro algebra with central charge c and highest weight h . It is shown in [FZ] that $L(c, 0)$ has a structure of a simple VOA. Here we consider the first unitary Virasoro VOA $L(1/2, 0)$. It is proved in [DMZ] [W] that $L(1/2, 0)$ is a rational C_2 -cofinite VOA of CFT-type and has exactly three irreducible modules, $L(1/2, 0)$, $L(1/2, 1/2)$ and $L(1/2, 1/16)$. Their fusion rules have also been computed and are as follows:

$$\begin{aligned}
L(1/2, 1/2) \times L(1/2, 1/2) &= L(1/2, 0), \\
L(1/2, 1/2) \times L(1/2, 1/16) &= L(1/2, 1/16), \\
L(1/2, 1/16) \times L(1/2, 1/16) &= L(1/2, 0) + L(1/2, 1/2).
\end{aligned} \tag{3.1}$$

First, we present an explicit realization of $L(1/2, 0)$ and its modules.

3.1 Ising model

In this section we give an explicit construction of the Ising model SVOA $L(1/2, 0) \oplus L(1/2, 1/2)$ and its \mathbb{Z}_2 -twisted modules $L(1/2, 1/16)^\pm$. This construction is well-known and the most of contents in this section can be found in [FFR], [FRW] and [KR].

Let \mathcal{A}_ψ be the algebra generated by $\{\psi_k \mid k \in \mathbb{Z} + \frac{1}{2}\}$ subject to the defining relation

$$[\psi_m, \psi_n]_+ := \psi_m \psi_n + \psi_n \psi_m = \delta_{m+n, 0}, \quad m, n \in \mathbb{Z} + \frac{1}{2},$$

and denote a subalgebra of \mathcal{A}_ψ generated by $\{\psi_k \mid k \in \mathbb{Z} + \frac{1}{2}, k > 0\}$ by \mathcal{A}_ψ^+ . Let $\mathbb{C}\mathbf{1}$ be a trivial \mathcal{A}_ψ^+ -module. Define a canonical induced \mathcal{A}_ψ -module M by

$$M := \text{Ind}_{\mathcal{A}_\psi^+}^{\mathcal{A}_\psi} \mathbb{C}\mathbf{1} = \mathcal{A}_\psi \otimes_{\mathcal{A}_\psi^+} \mathbb{C}\mathbf{1}.$$

Consider the generating series

$$\psi(z) := \sum_{n \in \mathbb{Z}} \psi_{n+\frac{1}{2}} z^{-n-1}.$$

Since $[\psi(z), \psi(w)]_+ = z^{-1} \delta(\frac{w}{z})$, $\psi(z)$ is local with itself. So we can consider a subalgebra of a local system on M generated by $\psi(z)$ and $I(z) = \text{id}_M$. By a direct calculation, one sees that the component operators of the generating series

$$\frac{1}{2} \psi(z) \circ_{-2} \psi(z) := \frac{1}{2} \text{Res}_{z_0} \{ (z_0 - z)^{-2} \psi(z_0) \psi(z) + (-z + z_0)^{-2} \psi(z_0) \psi(z) \}$$

defines a representation of the Virasoro algebra on M with central charge $1/2$, where \circ_n denotes the n -th normal ordered product defined in [Li1]. It follows from [KR] that M as a Vir-module is isomorphic to $L(1/2, 0) \oplus L(1/2, 1/2)$. Therefore, there is a unique simple vertex operator superalgebra structure on M such that $Y_M(\psi_{-\frac{1}{2}} \mathbf{1}, z) = \psi(z)$.

Theorem 3.1. *On M , there is a unique simple vertex operator superalgebra structure $(M, Y_M(\cdot, z), \mathbf{1}, \frac{1}{2} \psi_{-\frac{3}{2}} \psi_{-\frac{1}{2}} \mathbf{1})$ such that $Y_M(\psi_{-\frac{1}{2}} \mathbf{1}, z) = \psi(z)$.*

Another unitary Vir-module $L(1/2, 1/16)$ is realized as follows. Let \mathcal{A}_ϕ be the algebra generated by $\{\phi_n \mid n \in \mathbb{Z}\}$ with defining relation

$$[\phi_m, \phi_n]_+ = \delta_{m+n, 0}, \quad m, n \in \mathbb{Z}.$$

Let \mathcal{A}_ϕ^+ be a subalgebra of \mathcal{A}_ϕ generated by $\{\phi_n | n > 0\}$ and let $\mathbb{C}v_0$ be a trivial one-dimensional \mathcal{A}_ϕ^+ -module. Then set $N = \text{Ind}_{\mathcal{A}_\phi^+}^{\mathcal{A}_\phi} \mathbb{C}v_0$ as we did previously. We can find an action of the Virasoro algebra on N . Consider the generating series

$$\phi(z) := \sum_{n \in \mathbb{Z}} \phi_n z^{-n-\frac{1}{2}}.$$

By direct calculations one can show that $\phi(z)$ is local with itself. Consider a local system on N containing $\phi(z)$. Since the powers of z in $\phi(z)$ lie in $\mathbb{Z} + \frac{1}{2}$, we have to use the \mathbb{Z}_2 -twisted normal ordered product in [Li2]. Define a generating series $L(z)$ of operators on N by

$$\begin{aligned} L(z) &:= \phi(z) \circ_{-2} \phi(z) \\ &= \frac{1}{2} \text{Res}_{z_0} \text{Res}_{z_1} z_0^{-2} \left(\frac{z_1 - z_0}{z} \right)^{\frac{1}{2}} \\ &\quad \times \left\{ z_0^{-1} \delta \left(\frac{z_1 - z}{z_0} \right) \phi(z_1) \phi(z) + z_0^{-1} \delta \left(\frac{-z + z_1}{z_0} \right) \phi(z) \phi(z_1) \right\}, \end{aligned}$$

where \circ_n above denotes the n -th normal ordered product in a \mathbb{Z}_2 -twisted local system on N (cf. [Li2]). Then by a direct computation we find that the component operators of $L(z)$ defines a representation of the Virasoro algebra on N with central charge $1/2$. Set $v_{1/16}^\pm := \phi_0 \mathbb{1} \pm (1/\sqrt{2}) \mathbb{1}$. Then $v_{1/16}^\pm$ are highest weight vectors for the Virasoro algebra and the following decomposition is shown in [KR]:

$$N = L(1/2, 1/16)^+ \oplus L(1/2, 1/16)^-,$$

where $L(1/2, 1/16)^\pm$ are highest weight Vir-module generated by $v_{1/16}^\pm$, respectively.

Theorem 3.2. *The following \mathbb{Z}_2 -twisted Jacobi identity holds on N :*

$$\begin{aligned} & z_0^{-1} \delta \left(\frac{z_1 - z_2}{z_0} \right) Y_N(a, z_1) Y_N(b, z_2) - (-1)^{\varepsilon(a,b)} z_0^{-1} \left(\frac{-z_2 + z_1}{z_0} \right) Y_N(b, z_2) Y_N(a, z_1) \\ &= z_1^{-1} \left(\frac{z_2 + z_0}{z_1} \right) \left(\frac{z_2 + z_0}{z_1} \right)^{\varepsilon(a,a)/2} Y_N(Y_M(a, z_0) b, z_2), \end{aligned}$$

where $a, b \in M = L(1/2, 0) \oplus L(1/2, 1/2)$ and $\varepsilon(\cdot, \cdot)$ denotes the standard parity function. Therefore, the vertex operator map $Y_N(\cdot, z)$ defines inequivalent irreducible \mathbb{Z}_2 -twisted $L(1/2, 0) \oplus L(1/2, 1/2)$ -module structures on $L(1/2, 1/16)^\pm$.

3.2 Miyamoto involution

Let $(V, Y_V(\cdot, z), \mathbb{1}, \omega)$ be a VOA. A vector $e \in V$ is called a *conformal vector* if its component operators $Y_V(e, z) = \sum_{n \in \mathbb{Z}} e_{(n)} z^{-n-1} = \sum_{n \in \mathbb{Z}} L^e(n) z^{-n-2}$ generate a representation

of the Virasoro algebra on V :

$$[L^e(m), L^e(n)] = (m - n)L^e(m + n) + \delta_{m+n,0} \frac{m^3 - m}{12} c_e.$$

The scalar c_e is called *central charge* of a conformal vector e . We denote by $\text{Vir}(e)$ the sub VOA generated by e . If $\text{Vir}(e)$ is a rational VOA, then e is called a *rational conformal vector*. A decomposition $\omega = e + (\omega - e)$ is called *orthogonal* if both e and $\omega - e$ are conformal vectors and their component operators are mutually commutative.

Now assume that $e \in V$ is a rational conformal vector with central charge $1/2$. Then $\text{Vir}(e)$ is isomorphic to $L(1/2, 0)$. Since $L(1/2, 0)$ is rational, we can decompose V into a direct sum of irreducible $\text{Vir}(e)$ -modules as follows:

$$V = V_e(0) \oplus V_e(1/2) \oplus V_e(1/16),$$

where $V_e(h)$, $h \in \{0, 1/2, 1/16\}$, denotes the sum of all irreducible $\text{Vir}(e)$ -submodules of V isomorphic to $L(1/2, h)$. By the fusion rules (3.1), we have the following grading structure (cf. [M1]):

$$\begin{aligned} V_e(0) \cdot V_e(h) &\subset V_e(h), \quad h = 0, 1/2, 1/16, & V_e(1/2) \cdot V_e(1/2) &\subset V_e(0), \\ V_e(1/2) \cdot V_e(1/16) &\subset V_e(1/16), & V_e(1/16) \cdot V_e(1/16) &\subset V_e(0) \oplus V_e(1/2). \end{aligned}$$

Therefore, if $V_e(1/16) \neq 0$, then the linear map

$$\tau_e := 1 \quad \text{on } V_e(0) \oplus V_e(1/2), \quad -1 \quad \text{on } V_e(1/16)$$

defines an involutive automorphism on V (cf. [M1]). We call τ_e the *first Miyamoto involution* or simply *Miyamoto involution* associated to a conformal vector e . If $V_e(1/16) = 0$, then we can also define another involution as follows (cf. [M1]):

$$\sigma_e := 1 \quad \text{on } V_e(0), \quad -1 \quad \text{on } V_e(1/2).$$

We call σ_e the *second Miyamoto involution* associated to e .

Remark 3.3. It is shown in [C] and [M1] that the Miyamoto involution τ_e associated to a conformal vector e of the moonshine VOA [FLM] with central charge $1/2$ defines a 2A-involution of the Monster.

3.3 Commutant superalgebra

We keep the same notation as in the previous subsection. Let V be a simple VOA of CFT-type and $e \in V$ a rational conformal vector with central charge $1/2$. Set $T_e(h) := \{v \in V \mid L^e(0)v = h \cdot v\}$ for $h = 0, 1/2, 1/16$. Then $T_e(h)$ is a space of highest weight vectors

for $\text{Vir}(e)$ and is canonically isomorphic to $\text{Hom}_{\text{Vir}(e)}(L(1/2, h), V)$ for $h = 0, 1/2, 1/16$. Therefore, we have a decomposition as follows:

$$V = L(1/2, 0) \otimes T_e(0) \oplus L(1/2, 1/2) \otimes T_e(1/2) \oplus L(1/2, 1/16) \otimes T_e(1/16).$$

Lemma 3.4. *A decomposition $\omega = e + (\omega - e)$ is orthogonal.*

Proof: We compute $e_{(1)}\omega_{(2)}e$.

$$\begin{aligned} e_{(1)}\omega_{(2)}e &= \omega_{(2)}e_{(1)}e + [e_{(1)}, \omega_{(2)}]e \\ &= 2\omega_{(2)}e - [\omega_{(2)}, e_{(1)}]e \\ &= 2\omega_{(2)}e - \{(\omega_{(0)}e)_{(3)} + 2(\omega_{(1)}e)_{(2)} + (\omega_{(2)}e)_{(1)}\}e \\ &= 2\omega_{(2)}e - (\omega_{(2)}e)_{(1)}e. \end{aligned}$$

By the skew-symmetry, we have $(\omega_{(2)}e)_{(1)}e = e_{(1)}\omega_{(2)}e - \omega_{(0)}e_{(2)}\omega_{(2)}e$. Since $e_{(2)}\omega_{(2)}e \in V_0 = \mathbb{C}\mathbf{1}$, $\omega_{(0)}e_{(2)}\omega_{(2)}e = 0$ and so $(\omega_{(2)}e)_{(1)}e = e_{(1)}\omega_{(2)}e$. Substituting this into the equality above, we get $e_{(1)}\omega_{(2)}e = \omega_{(2)}e$. Namely, $\omega_{(2)}e$ is an eigenvector for $e_{(1)}$ with eigenvalue 1. Since V is a module for $\text{Vir}(e)$, there is no eigenvector with $e_{(1)}$ -weight 1. Hence $\omega_{(2)}e = 0$. Then the assertion follows from Theorem 5.1 of [FZ]. \blacksquare

Recall the commutant subalgebra $\text{Com}_V(\text{Vir}(e)) := \ker_V e_{(0)}$ defined in [FZ]. By the lemma above, $(T_e(0), \omega - e)$ forms a sub VOA of V whose action on V is commutative with that of $\text{Vir}(e)$ on V . In particular, $T_e(h)$, $h = 0, 1/2, 1/16$, are $T_e(0)$ -modules.

Proposition 3.5. (1) $T_e(0) = \ker_V e_{(0)} = \text{Com}_V(\text{Vir}(e))$ is a simple sub VOA with the Virasoro vector $\omega - e$.

(2) $T_e(1/2)$ is an irreducible $T_e(0)$ -module.

(3) $\text{Vir}(e) = \ker_V(\omega - e)_{(0)} = \text{Com}_V(T_e(0))$.

Proof: (1): Let $v \in V$. Since $e_{(1)}v = 0$ implies $e_{(0)}v = 0$, $T_e(0) = \ker_V e_{(1)} = \ker_V e_{(0)}$. So we only need to show that $T_e(0)$ is simple. Since V is simple, the τ_e -orbifold $V^{\langle \tau_e \rangle} = V_e(0) \oplus V_e(1/2)$ is simple. Then the σ_e -orbifold $(V^{\langle \tau_e \rangle})^{\langle \sigma_e \rangle} = V_e(0)$ is also simple. Since $\text{Vir}(e) \otimes T_e(0) \ni a \otimes b \mapsto a_{(-1)}b \in V_e(0)$ is an isomorphism of VOAs, $T_e(0)$ is also simple.

(2): Since both $V^{\langle \tau_e \rangle} = V_e(0) \oplus V_e(1/2)$ and $V_e(0)$ are simple VOAs, $V_e(1/2)$ is an irreducible $V_e(0)$ -module. So $T_e(1/2)$ is also irreducible.

(3): As $\omega - e$ is a conformal vector, $\ker_V(\omega - e)_{(0)}$ is generally contained in $\ker_V(\omega - e)_{(1)}$. On the other hand, since V is of CFT-type, $\ker_V(\omega - e)_{(1)} = \text{Vir}(e)$. Then

$$\text{Vir}(e) \subset \text{Com}_V(\text{Com}_V(\text{Vir}(e))) = \ker_V(\omega - e)_{(0)}$$

implies $\text{Vir}(e) = \ker_V(\omega - e)_{(0)}$. \blacksquare

Theorem 3.6. *Suppose that $T_e(1/2) \neq 0$. Then there exists a simple SVOA structure on $T_e(0) \oplus T_e(1/2)$ such that the even part of a tensor product of SVOAs*

$$\{L(1/2, 0) \oplus L(1/2, 1/2)\} \otimes \{T_e(0) \oplus T_e(1/2)\}$$

is isomorphic to $V_e(0) \oplus V_e(1/2)$ as a VOA.

Proof: We shall define vertex operators on an abstract space $T_e(0) \oplus T_e(1/2)$. First, we show an existence of a $T_e(0)$ -intertwining operator of type $T_e(1/2) \times T_e(1/2) \rightarrow T_e(0)$. Write $Y_V(\omega, z) = \sum_{n \in \mathbb{Z}} L(n)z^{-n-2}$ and $Y_V(e, z) = \sum_{n \in \mathbb{Z}} L^e(n)z^{-n-2}$. Since $L(0) - L^e(0)$ semisimply acts on both $T_e(0)$ and $T_e(1/2)$, we can take bases $\{a^\gamma \mid \gamma \in \Gamma\}$ and $\{u^\lambda \mid \lambda \in \Lambda\}$ of $T_e(0)$ and $T_e(1/2)$, respectively, consisting of eigenvectors for $L(0) - L^e(0)$. Let $\pi_\gamma : V_e(0) \rightarrow L(1/2, 0) \otimes a^\gamma$, $\gamma \in \Gamma$, be a projection map. For $\gamma \in \Gamma$ and $\lambda, \mu \in \Lambda$, we define a linear operator $I_{\lambda\mu}^\gamma(\cdot, z)$ of type $L(1/2, 1/2) \times L(1/2, 1/2) \rightarrow L(1/2, 0) \otimes a^\gamma$ by

$$\begin{aligned} I_{\lambda\mu}^\gamma(x, z)y &:= z^{-L(0)+L^e(0)}\pi_\gamma Y(z^{L(0)-L^e(0)}x \otimes u^\lambda, z)z^{L(0)-L^e(0)}y \otimes u^\mu \\ &= z^{-|\gamma|+|\lambda|+|\mu|}\pi_\gamma Y(x \otimes u^\lambda, z)y \otimes u^\mu, \end{aligned}$$

for $x, y \in L(1/2, 1/2)$, where $|\lambda|$, $|\mu|$ and $|\gamma|$ denote the $(L(0) - L^e(0))$ -weight of u^λ , u^μ and a^γ , respectively. Then by [DL] [M1] the operator $I_{\lambda\mu}^\gamma(\cdot, z)$ is an $L(1/2, 0)$ -intertwining operator of type $L(1/2, 1/2) \times L(1/2, 1/2) \rightarrow L(1/2, 0)$. Since the space of intertwining operators of that type is one-dimensional, each $I_{\lambda\mu}^\gamma(\cdot, z)$ is proportional to the vertex operator map $Y_M(\cdot, z)$ on the SVOA $M = L(1/2, 0) \oplus L(1/2, 1/2)$ which we constructed explicitly in Section 3.1. Thus there exist scalars $c_{\lambda\mu}^\gamma \in \mathbb{C}$ such that $I_{\lambda\mu}^\gamma(\cdot, z) = c_{\lambda\mu}^\gamma Y_M(\cdot, z)$. Then the vertex operator of $x \otimes u^\lambda \in L(1/2, 1/2) \otimes T_e(1/2)$ on $V_e(1/2)$ can be written as follows:

$$Y_V(x \otimes u^\lambda, z)y \otimes u^\mu = Y_M(x, z)y \otimes \sum_{\gamma \in \Gamma} c_{\lambda\mu}^\gamma a^\gamma z^{|\gamma|-|\lambda|-|\mu|}.$$

Thus, by setting $J(u^\lambda, z)u^\mu := \sum_{\gamma \in \Gamma} c_{\lambda\mu}^\gamma a^\gamma z^{|\gamma|-|\lambda|-|\mu|}$, we obtain a decomposition

$$Y_V(x \otimes u^\lambda, z)y \otimes u^\mu = Y_M(x, z)y \otimes J(u^\lambda, z)u^\mu$$

for $x \otimes u^\lambda, y \otimes u^\mu \in L(1/2, 1/2) \otimes V_e(1/2)$. We claim that $J(\cdot, z)$ is a $T_e(0)$ -intertwining operator of type $T_e(1/2) \times T_e(1/2) \rightarrow T_e(0)$. It is obvious that $J(u, z)v$ contains finitely many negative powers of z and the $(\omega - e)_{(0)}$ -derivation property $J((\omega - e)_0 u, z)v = \frac{d}{dz}J(u, z)v$ hold for all $u, v \in T_e(1/2)$. So we should show that $J(\cdot, z)$ satisfies both the commutativity and the associativity. Let $a \in T_e(0)$ and $u, v \in T_e(1/2)$ be arbitrary elements. Then the commutativity of vertex operators on V gives

$$\begin{aligned} (z_1 - z_2)^N Y_V(\mathbf{1} \otimes a, z_1) Y_V(\psi_{-\frac{1}{2}} \mathbf{1} \otimes u, z_2) \psi_{-\frac{1}{2}} \mathbf{1} \otimes v \\ = (z_1 - z_2)^N Y_V(\psi_{-\frac{1}{2}} \mathbf{1} \otimes u, z_2) Y_V(\mathbf{1} \otimes a, z_1) \psi_{-\frac{1}{2}} \mathbf{1} \otimes v \end{aligned}$$

for sufficiently large N . Rewriting the equality above we get

$$\begin{aligned} & (z_1 - z_2)^N Y_M(\psi_{-\frac{1}{2}} \mathbf{1}, z_2) \psi_{-\frac{1}{2}} \mathbf{1} \otimes Y_{T_e(0)}(a, z_1) J(u, z_2) v \\ &= (z_1 - z_2)^N Y_M(\psi_{-\frac{1}{2}} \mathbf{1}, z_2) \psi_{-\frac{1}{2}} \mathbf{1} \otimes J(u, z_2) Y_{T_e(1/2)}(a, z_1) v, \end{aligned}$$

where $Y_{T_e(0)}(a, z)$ and $Y_{T_e(1/2)}(\cdot, z)$ denote the vertex operator of $a \in T_e(0)$ on $T_e(0)$ and $T_e(1/2)$, respectively. By comparing the coefficients of $(\psi_{-\frac{1}{2}} \mathbf{1})_{(0)} \psi_{-\frac{1}{2}} \mathbf{1} = \mathbf{1}$, we get the commutativity:

$$(z_1 - z_2)^N Y_{T_e(0)}(a, z_1) J(u, z_2) v = (z_1 - z_2)^N J(u, z_2) Y_{T_e(1/2)}(a, z_1) v.$$

Similarly, by considering coefficients of $Y_V(Y_V(\mathbf{1} \otimes a, z_0) \psi_{-\frac{1}{2}} \mathbf{1} \otimes u, z_2) \psi_{-\frac{1}{2}} \mathbf{1} \otimes v$ in V , we obtain the associativity:

$$(z_0 + z_2)^N Y_{T_e(0)}(a, z_0 + z_2) J(u, z_2) v = (z_2 + z_0)^N J(Y_{T_e(1/2)}(a, z_0) u, z_2) v.$$

Hence, $J(\cdot, z)$ is a $T_e(0)$ -intertwining operator of the desired type.

Using $Y_V(\cdot, z)$ and $J(\cdot, z)$, we introduce a vertex operator map $\hat{Y}(\cdot, z)$ on $T_e(0) \oplus T_e(1/2)$. Let $a, b \in T_e(0)$ and $u, v \in T_e(1/2)$. We define

$$\begin{aligned} \mathbf{1} \otimes \hat{Y}(a, z) b &:= Y_V(\mathbf{1} \otimes a, z) \mathbf{1} \otimes b, \quad \psi_{-\frac{1}{2}} \mathbf{1} \otimes \hat{Y}(a, z) u := Y_V(\mathbf{1} \otimes a, z) \psi_{-\frac{1}{2}} \mathbf{1} \otimes u, \\ \psi_{-\frac{1}{2}} \mathbf{1} \otimes \hat{Y}(u, z) a &:= e^{z(L(-1) - L^e(-1))} Y_V(\mathbf{1} \otimes a, z) \psi_{-\frac{1}{2}} \mathbf{1} \otimes u, \quad \hat{Y}(u, z) v := J(u, z) v. \end{aligned}$$

Then all $\hat{Y}(\cdot, z)$ are $T_e(0)$ -intertwining operators. We note that $\hat{Y}(\cdot, z)$ satisfies the vacuum condition:

$$\hat{Y}(x, z) \mathbf{1} \in x + (T_e(0) \oplus T_e(1/2)) [[z]] z$$

for any $x \in T_e(0) \oplus T_e(1/2)$. Hence, to prove that $T_e(0) \oplus T_e(1/2)$ is a simple SVOA, it is sufficient to show that the vertex operator map $\hat{Y}(\cdot, z)$ defined above satisfies the commutativity. By our definition, the vertex operator map $Y_V(a \otimes x, z)$ of $a \otimes x \in L(1/2, h) \otimes T_e(h) = V_e(h)$, $h = 0, 1/2$, can be written as $Y_M(a, z) \otimes \hat{Y}(x, z)$. Because of our manifest construction of $Y_M(\cdot, z)$ in Section 3.1, we can perform explicit computations of the vertex operator $Y_M(\cdot, z)$ on $L(1/2, 0) \oplus L(1/2, 1/2)$. Therefore, by comparing the coefficients of vertex operators on V , we can prove that $\hat{Y}(\cdot, z)$ satisfies the (super-)commutativity. Thus, by our definition, $(T_e(0) \oplus T_e(1/2), \hat{Y}(\cdot, z), \mathbf{1}, \omega - e)$ carries a structure of a simple SVOA. The rest of the assertion is now clear. \blacksquare

Remark 3.7. There is another proof of Theorem 3.6 in [Hö1]. In [Hö1], he assumed the existence of a positive definite invariant bilinear form on a real form of V . However, our argument does not need the assumption on the unitary form.

Since $\tau_e^2 = 1$ on V , the space $V_e(1/16)$ is an irreducible $V^{\langle \tau_e \rangle}$ -module. As a $(V^{\langle \tau_e \rangle})^{\langle \sigma_e \rangle} = \text{Vir}(e) \otimes T_e(0)$ -module, $V_e(1/16)$ can be written as $L(1/2, 1/16) \otimes T_e(1/16)$. It is not clear that $T_e(1/16)$ is irreducible under $T_e(0)$. However, we can prove that it is irreducible under $T_e(0) \oplus T_e(1/2)$.

Theorem 3.8. *Suppose that $V_e(1/2) \neq 0$ and $V_e(1/16) \neq 0$. Then $T_e(1/16)$ carries a structure of an irreducible \mathbb{Z}_2 -twisted $T_e(0) \oplus T_e(1/2)$ -module. Moreover, $V_e(1/16)$ is isomorphic to a tensor product of an irreducible \mathbb{Z}_2 -twisted $L(1/2, 0) \oplus L(1/2, 1/2)$ -module $L(1/2, 1/16)^+$ and an irreducible \mathbb{Z}_2 -twisted $T_e(0) \oplus T_e(1/2)$ -module $T_e(1/16)$.*

Proof: The idea of the proof is the same as that of Theorem 3.6. Computing vertex operators on $L(1/2, 1/16)^+$ and then comparing the coefficients in V , we will reach the assertion. Denote by $Y_N(\cdot, z)$ the vertex operator map on the \mathbb{Z}_2 -twisted $L(1/2, 0) \oplus L(1/2, 1/2)$ -module $L(1/2, 1/16)^+$ as we constructed in Section 3.1. Let $a \otimes b \in L(1/2, h) \otimes T_e(h)$ with $h = 0$ or $1/2$ and $x \otimes y \in L(1/2, 1/16) \otimes T_e(1/16)$. As we did before, we can find $T_e(0)$ -intertwining operators $Y_{T_e(h) \times T_e(1/16)}(\cdot, z)$ of types $T_e(h) \times T_e(1/16) \rightarrow T_e(1/16)$ such that

$$Y_V(a \otimes b, z)x \otimes y = Y_N(a, z)x \otimes Y_{T_e(h) \times T_e(\frac{1}{16})}(b, z)y. \quad (3.2)$$

Define $\hat{Y}(b, z)y := Y_{T_e(h) \times T_e(\frac{1}{16})}(b, z)y$ for $b \in T_e(h)$, $h = 0, 1/2$ and $y \in T_e(1/16)$. By direct computations, we can prove that the \mathbb{Z}_2 -twisted Jacobi identity for $Y_N(\cdot, z)$ together with the Jacobi identity for $Y_V(\cdot, z)$ gives the \mathbb{Z}_2 -twisted Jacobi identity for $\hat{Y}(\cdot, z)$. Thus, $(T_e(1/16), \hat{Y}(\cdot, z))$ is a \mathbb{Z}_2 -twisted $T_e(0) \oplus T_e(1/2)$ -module. Since $V_e(1/16) = L(1/2, 1/16) \otimes T_e(1/16)$ is irreducible under $V_e(0) \oplus V_e(1/2)$, the irreducibility of $T_e(1/16)$ is obvious. ■

3.4 One point stabilizer

In the rest of this section we will work the following setup:

Hypothesis 1.

- (1) V is a holomorphic VOA of CFT-type.
- (2) e is a rational conformal vector of V with central charge $1/2$.
- (3) $V_e(h) \neq 0$ for $h = 0, 1/2, 1/16$.
- (4) $V_e(0)$ and $T_e(0)$ are rational C_2 -cofinite VOAs of CFT-type.
- (5) $V_e(1/16)$ is a simple current $V^{\langle \tau_e \rangle} = V_e(0) \oplus V_e(1/2)$ -module.
- (6) $T_e(1/2)$ is a simple current $T_e(0)$ -module.

Define the one-point stabilizer by $C_{\text{Aut}(V)}(e) := \{\rho \in \text{Aut}(V) \mid \rho(e) = e\}$. Then by $\tau_{\rho(e)} = \rho \tau_e \rho^{-1}$ for any $\rho \in \text{Aut}(V)$, we have $C_{\text{Aut}(V)}(e) \subset C_{\text{Aut}(V)}(\tau_e)$, where $C_{\text{Aut}(V)}(\tau_e)$ denotes the centralizer of an involution $\tau_e \in \text{Aut}(V)$.

Lemma 3.9. *There are group homomorphisms $\psi_1 : C_{\text{Aut}(V)}(e) \rightarrow C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e)$ and $\psi_2 : C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e) \rightarrow \text{Aut}(T_e(0))$ such that $\ker(\psi_1) = \langle \tau_e \rangle$ and $\ker(\psi_2) = \langle \sigma_e \rangle$.*

Proof: Let $\rho \in C_{\text{Aut}(V)}(e)$. Then ρ preserves the space of highest weight vectors $T_e(h)$ where $h \in \{0, 1/2, 1/16\}$. Then we can define the actions of ρ on the space of highest weight vectors $T_e(h)$ and the components $V_e(h)$ for $h \in \{0, 1/2, 1/16\}$. In particular, we have group homomorphisms $\psi_1 : C_{\text{Aut}(V)}(e) \rightarrow C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e)$ and $\psi_2 : C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e) \rightarrow \text{Aut}(T_e(0))$ by a natural way. Assume that $\psi_1(\rho) = \text{id}_{V^{\langle \tau_e \rangle}}$ for $\rho \in C_{\text{Aut}(V)}(e)$. Since $\rho \in C_{\text{Aut}(V)}(\tau_e)$, ρ acts on $V_e(1/16)$ and commutes with the action of $V^{\langle \tau_e \rangle} = V_e(0) \oplus V_e(1/2)$ on its module $V_e(1/16)$. Therefore, ρ on $V_e(1/16)$ is a scalar by Schur's lemma and hence $\rho \in \langle \tau_e \rangle \subset C_{\text{Aut}(V)}(\tau_e)$. Similarly, if $\psi_1(\rho') = \text{id}_{T_e(0)}$ for $\rho' \in C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e)$, then $\rho' \in \langle \sigma_e \rangle \subset C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e)$. ■

Theorem 3.10. *Under Hypothesis 1, $V^{\langle \tau_e \rangle}$ has exactly four inequivalent irreducible modules, $V^{\langle \tau_e \rangle}$, $V_e(1/16)$, $W^0 := L(1/2, 0) \otimes T_e(1/2) \oplus L(1/2, 1/2) \otimes T_e(0)$ and*

$$W^1 := V_e(1/16) \boxtimes_{V^{\langle \tau_e \rangle}} W^0.$$

Proof: Note that $V_e(0) = \text{Vir}(e) \otimes T_e(0)$ and $V^{\langle \tau_e \rangle}$ are simple rational C_2 -cofinite VOAs of CFT-type under Hypothesis 1. Therefore, we can apply a theory of fusion products here. Since $V = V^{\langle \tau_e \rangle} \oplus V_e(1/16)$ is a \mathbb{Z}_2 -graded simple current extension of $V^{\langle \tau_e \rangle}$, every irreducible $V^{\langle \tau_e \rangle}$ -module is lifted to be either an irreducible V -module or an irreducible τ_e -twisted V -module. Moreover, the τ_e -twisted V -module is unique up to isomorphism by Theorem 10.3 of [DLM2]. Consider a $V_e(0)$ -module $L(1/2, 1/2) \otimes T_e(0)$. Since $T_e(1/2)$ is a simple current $T_e(0)$ -module, the space

$$W^0 = L(1/2, 1/2) \otimes T_e(0) \oplus L(1/2, 0) \otimes T_e(1/2)$$

has a unique structure of an irreducible $V^{\langle \tau_e \rangle}$ -module by Theorem 2.11. We note that the top weight of W^0 is half-integral. Thus the induced module

$$W = W^0 \oplus W^1, \quad W^1 = V_e(1/16) \boxtimes_{V^{\langle \tau_e \rangle}} W^0,$$

becomes an irreducible τ_e -twisted V -module again by Theorem 2.11. Therefore, $V^{\langle \tau_e \rangle}$ has exactly four irreducible modules as in the assertion. Finally we remark that $V^{\langle \tau_e \rangle}$, $V_e(1/16)$ and W^1 have integral top weights. ■

By the fusion rules (3.1), we note that W^1 as a $\text{Vir}(e)$ -module is a direct sum of copies of $L(1/2, 1/16)$. Set the space of highest weight vectors of W^1 by $Q_e(1/16) := \{v \in W^1 \mid L^e(0)v = (1/16) \cdot v\}$. Then as a $\text{Vir}(e) \otimes T_e(0)$ -module, $W^1 \simeq L(1/2, 1/16) \otimes Q_e(1/16)$. In this case, we can also verify that the space $Q_e(1/16)$ naturally carries an irreducible \mathbb{Z}_2 -twisted $T_e(0) \oplus T_e(1/2)$ -module structure.

Proposition 3.11. *If the \mathbb{Z}_2 -twisted $T_e(0) \oplus T_e(1/2)$ -module $T_e(1/16)$ is irreducible as a $T_e(0)$ -module, then its \mathbb{Z}_2 -conjugate is isomorphic to $Q_e(1/16)$ as a \mathbb{Z}_2 -twisted $T_e(0) \oplus T_e(1/2)$ -module. In this case there are three irreducible $T_e(0)$ -modules, $T_e(0)$, $T_e(1/2)$ and $T_e(1/16)$. Conversely, if $T_e(1/16)$ as a $T_e(0)$ -module is not irreducible, then so is $Q_e(1/16)$ and in this case there are six inequivalent irreducible $T_e(0)$ -modules.*

Proof: Assume that $T_e(1/16)$ is irreducible as a $T_e(0)$ -module. Then its \mathbb{Z}_2 -conjugate is not isomorphic to $T_e(1/16)$ as a \mathbb{Z}_2 -twisted $T_e(0) \oplus T_e(1/2)$ -module. We denote the \mathbb{Z}_2 -conjugate of $T_e(1/16)$ by $T_e(1/16)^-$. It is shown in Theorem 2.12 that every irreducible $T_e(0)$ -module is lifted to be either an irreducible $T_e(0) \oplus T_e(1/2)$ -module or an irreducible \mathbb{Z}_2 -twisted $T_e(0) \oplus T_e(1/2)$ -module. Then by the classification of irreducible $V^{\langle \tau_e \rangle}$ -modules in Theorem 3.10, we see that any \mathbb{Z}_2 -twisted irreducible $T_e(0) \oplus T_e(1/2)$ -module is isomorphic to one and only one of $T_e(1/16)$ and $T_e(1/16)^- = Q_e(1/16)$.

Conversely, if $T_e(1/16)$ is not irreducible, then it is a direct sum of two inequivalent irreducible $T_e(0)$ -module as $T_e(1/2)$ is a simple current $T_e(0)$ -module. Then $Q_e(1/16)$ is also a direct sum of two inequivalent irreducible $T_e(0)$ -modules and $Q_e(1/16) \not\cong T_e(1/16)$ as $T_e(0)$ -modules because of the classification of irreducible $V^{\langle \tau_e \rangle}$ -modules. ■

Corollary 3.12. *If $T_e(1/16)$ is irreducible as a $T_e(0)$ -module, then $V^{\langle \tau_e \rangle} \oplus W^1$ is a \mathbb{Z}_2 -graded simple current extension of $V^{\langle \tau_e \rangle}$ which is isomorphic to $V = V^{\langle \tau_e \rangle} \oplus V_e(1/16)$.*

Proof: If $T_e(1/16)$ is an irreducible $T_e(0)$ -module, then by the previous proposition the σ_e -conjugate $V_e(0) \oplus V_e(1/2)$ -module of $V_e(1/16) = L(1/2, 1/16) \otimes T_e(1/16)$ is isomorphic to $W^1 = L(1/2, 1/16) \otimes Q_e(1/16)$. Then as the σ_e -conjugate extension of $V^{\langle \tau_e \rangle} \oplus V_e(1/16)$, $V^{\langle \tau_e \rangle} \oplus W^1$ has a structure of a \mathbb{Z}_2 -graded extension. ■

Remark 3.13. The above corollary implies that the \mathbb{Z}_2 -twisted orbifold construction applied to V in the case of $\mathbb{Z}_2 = \langle \tau_e \rangle$ yields again V itself.

Theorem 3.14. *Under Hypothesis 1,*

- (i) ψ_2 is surjective, that is, $C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e) \simeq \langle \sigma_e \rangle \cdot \text{Aut}(T_e(0))$.
- (ii) $\text{Aut}(T_e(0) \oplus T_e(1/2)) \simeq 2 \cdot (C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e) / \langle \sigma_e \rangle)$, where 2 denotes the canonical \mathbb{Z}_2 -symmetry on the SVOA $T_e(0) \oplus T_e(1/2)$.
- (iii) $|C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e) : C_{\text{Aut}(V)}(e) / \langle \tau_e \rangle| \leq 2$.
- (iv) If $C_{\text{Aut}(V)}(e) / \langle \tau_e \rangle$ is simple or has an odd order, then extensions in (i) and (ii) split. That is, $C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e) \simeq \langle \sigma_e \rangle \times C_{\text{Aut}(V)}(e) / \langle \tau_e \rangle$ and $\text{Aut}(T_e(0) \oplus T_e(1/2)) \simeq 2 \times \text{Aut}(T_e(0))$.

Proof: We have an injection from $C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e) / \langle \sigma_e \rangle$ to $\text{Aut}(T_e(0))$ by Lemma 3.9. We show that every element in $\text{Aut}(T_e(0))$ lifts to be an element in $C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e)$. By Proposition 3.11, every irreducible $T_e(0)$ -module appears in one of $T_e(0)$, $T_e(1/2)$, $T_e(1/16)$

or $Q_e(1/16)$ as a submodule. In particular, we find that $T_e(0)$ is the only irreducible $T_e(0)$ -module whose top weight is integral and $T_e(1/2)$ is the only irreducible $T_e(0)$ -module whose top weight is in $1/2 + \mathbb{N}$. Let $\rho \in \text{Aut}(T_e(0))$. Then by considering top weights we can immediately see that $T_e(0)^\rho \simeq T_e(0)$ and $T_e(1/2)^\rho \simeq T_e(1/2)$. Then by Theorem 2.16 we have a lifting $\tilde{\rho} \in \text{Aut}(T_e(0) \oplus T_e(1/2))$ such that $\tilde{\rho}T_e(0) = T_e(0)$, $\tilde{\rho}T_e(1/2) = T_e(1/2)$ and $\tilde{\rho}|_{T_e(0)} = \rho$. Since this lifting is unique up to a multiple of the canonical \mathbb{Z}_2 -symmetry on $T_e(0) \oplus T_e(1/2)$, we have $\text{Aut}(T_e(0) \oplus T_e(1/2)) \simeq 2 \cdot \text{Aut}(T_e(0))$. Now consider the canonical extension of $\tilde{\rho}$ to $C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e)$. We define $\tilde{\rho} \in C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e)$ by

$$\tilde{\rho}|_{L(1/2, h) \otimes T_e(h)} = \text{id}_{L(1/2, h)} \otimes \tilde{\rho}$$

for $h = 0, 1/2$. Then by this lifting $C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e)$ contains a subgroup isomorphic to $2 \cdot \text{Aut}(T_e(0))$. Moreover, the canonical \mathbb{Z}_2 -symmetry on the SVOA $T_e(0) \oplus T_e(1/2)$ is naturally extended to $\sigma_e \in C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e)$. Clearly $\psi_2(\tilde{\rho}) = \rho$ and so ψ_2 is surjective. Hence we have the desired isomorphisms $C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e) \simeq \langle \sigma_e \rangle \cdot \text{Aut}(T_e(0))$ and $\text{Aut}(T_e(0) \oplus T_e(1/2)) \simeq 2 \cdot (C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e) / \langle \sigma_e \rangle)$. This completes the proof of (i) and (ii).

Consider (iii). By Theorem 3.10, there are exactly three irreducible $V^{\langle \tau_e \rangle}$ -modules whose top weights are integral, namely, $V^{\langle \tau_e \rangle}$, $V_e(1/16)$ and W^1 . Since $C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e)$ acts on the 2-point set $\{V_e(1/16), W^1\}$ as a permutation, there is a subgroup H of $C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e)$ of index at most 2 such that $V_e(1/16)^\pi \simeq V_e(1/16)$ as a $V^{\langle \tau_e \rangle}$ -module for all $\pi \in H$. Then there is a lifting $\tilde{\pi} \in C_{\text{Aut}(V)}(e)$ of π such that $\psi_1(\tilde{\pi}) = \pi$ for each $\pi \in H$ by Theorem 2.16. Thus $|C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e) : C_{\text{Aut}(V)}(e) / \langle \tau_e \rangle| \leq 2$ and (iii) holds.

Consider (iv). Suppose that $C_{\text{Aut}(V)}(e) / \langle \tau_e \rangle$ is a simple group or an odd group. Then $C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e)$ contains a simple group $C_{\text{Aut}(V)}(e) / \langle \tau_e \rangle$ with index at most 2 by (iii). However, since $C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e)$ contains a normal subgroup $\langle \sigma_e \rangle$ of order 2, the index $|C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e) : C_{\text{Aut}(V)}(e) / \langle \tau_e \rangle|$ must be 2 and hence we obtain the desired isomorphism $C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e) = \langle \sigma_e \rangle \times C_{\text{Aut}(V)}(e) / \langle \tau_e \rangle$. In this case, it is easy to see that the extension $\text{Aut}(T_e(0) \oplus T_e(1/2)) = 2 \cdot \text{Aut}(T_e(0))$ splits. ■

Corollary 3.15. *If $C_{\text{Aut}(V)}(e) / \langle \tau_e \rangle$ is simple, then $V_e(1/16)$ is an irreducible $V_e(0)$ -module and $T_e(1/16)$ is an irreducible $T_e(0)$ -module. Therefore, $V^{\langle \tau_e \rangle} \oplus W^1$ forms the σ_e -conjugate extension of $V = V^{\langle \tau_e \rangle} \oplus V_e(1/16)$ and is isomorphic to V .*

Proof: Let H be the subgroup of $C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e)$ which fixes $V_e(1/16)$ in the action on the 2-point set $\{V_e(1/16), W^1\}$. It is shown in the proof of (iii) of Theorem 3.14 that we have inclusions

$$H \subset C_{\text{Aut}(V)}(e) / \langle \tau_e \rangle \subset C_{\text{Aut}(V^{\langle \tau_e \rangle})}(e) = \langle \sigma_e \rangle \times C_{\text{Aut}(V)}(e) / \langle \tau_e \rangle.$$

Therefore, $\sigma_e \notin H$ and hence the σ_e permutes $V_e(1/16)$ and W^1 . Then $V_e(1/16)$ is an irreducible $V_e(0)$ -module by Proposition 3.11 and hence $T_e(1/16)$ as a $T_e(0)$ -module is irreducible. The rest of the assertion is now clear. ■

Remark 3.16. A result similar to the assertion (iii) of Theorem 3.14 is already established in [M6]. Also, we should note that the idea of the above proof is already developed in [Sh].

4 2A-framed VOA

Definition 4.1. A simple vertex operator algebra (V, ω) is called *2A-framed* if there is an orthogonal decomposition $\omega = e^1 + \cdots + e^n$ such that each e^i generates a sub VOA isomorphic to $L(1/2, 0)$. The decomposition $\omega = e^1 + \cdots + e^n$ is called a *2A-frame* of V .

Remark 4.2. As shown in [DMZ], the Leech lattice VOA V_Λ and the moonshine VOA V^\natural are examples of 2A-framed VOAs.

Let (V, ω) be a 2A-framed VOA with a 2A-frame $\omega = e^1 + \cdots + e^n$. Set $T := \text{Vir}(e^1) \otimes \cdots \otimes \text{Vir}(e^n)$, where $\text{Vir}(e^i)$ denotes the sub VOA generated by e^i . Then $T \simeq L(1/2, 0)^{\otimes n}$ and V is a direct sum of irreducible T -submodules $\otimes_{i=1}^n L(1/2, h_i)$ with $h_i \in \{0, 1/2, 1/16\}$. For each irreducible T -module $\otimes_{i=1}^n L(1/2, h_i)$, we associate its 1/16-word $(\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_2^n$ by the rule $\alpha_i = 1$ if and only if $h_i = 1/16$. For each $\alpha \in \mathbb{Z}_2^n$, denote by V^α the sum of all irreducible T -submodules whose 1/16-words are equal to α and define a linear code $S \subset \mathbb{Z}_2^n$ by $S = \{\alpha \in \mathbb{Z}_2^n \mid V^\alpha \neq 0\}$. Then we have the *1/16-word decomposition* $V = \oplus_{\alpha \in S} V^\alpha$ of V . By the fusion rules (3.1), we have an S -graded structure $V^\alpha \cdot V^\beta \subset V^{\alpha+\beta}$. Namely, the dual group S^* of an abelian 2-group S acts on V , and we find that this automorphism group coincides with the elementary abelian 2-group generated by the first Miyamoto involutions $\{\tau_{e^i} \mid 1 \leq i \leq n\}$. Therefore, all V^α , $\alpha \in S$, are irreducible $V^{S^*} = V^0$ -modules by [DM1]. Since there is no $L(1/2, 1/16)$ -component in V^0 , the fixed point subalgebra $V^{S^*} = V^0$ has the following shape:

$$V^0 = \bigoplus_{h_i \in \{0, 1/2\}} m_{h_1, \dots, h_n} L(1/2, h_1) \otimes \cdots \otimes L(1/2, h_n),$$

where m_{h_1, \dots, h_n} denotes the multiplicity. On V^0 we can define the second Miyamoto involutions σ_{e^i} for $i = 1, \dots, n$. Denote by Q the elementary abelian 2-subgroup of $\text{Aut}(V^0)$ generated by $\{\sigma_{e^i} \mid 1 \leq i \leq n\}$. Then by the quantum Galois theory [DM1] we have $(V^0)^Q = T$ and each $m_{h_1, \dots, h_n} L(1/2, h_1) \otimes \cdots \otimes L(1/2, h_n)$ is an irreducible T -submodule. Thus $m_{h_1, \dots, h_n} \in \{0, 1\}$ and we obtain an even linear code $D := \{(2h_1, \dots, 2h_n) \in \mathbb{Z}_2^n \mid m_{h_1, \dots, h_n} \neq 0\}$ such that

$$V^0 = \bigoplus_{\alpha = (\alpha_1, \dots, \alpha_n) \in D} L(1/2, \alpha_1/2) \otimes \cdots \otimes L(1/2, \alpha_n/2). \quad (4.1)$$

We call a pair (D, S) the *structure codes* of a 2A-framed VOA V . Since powers of z in an $L(1/2, 0)$ -intertwining operator of type $L(1/2, 1/2) \times L(1/2, 1/2) \rightarrow L(1/2, 1/16)$ are

half-integral, structure codes satisfy $D \subset S^\perp$. Since all V^α , $\alpha \in S$, are irreducible modules for V^0 , the representation theory of V^0 is important to study a 2A-framed VOA V . We review Miyamoto's results on the code VOAs [M3] [M4] in the next subsection.

4.1 Code VOA

Below we often identify the code \mathbb{Z}_2^n with the power set of an n -point set $\Omega = \{1, 2, \dots, n\}$ with the symmetric difference operation. For $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_2^n$, we denote by $\text{Supp}(\alpha)$ the subset $\{i \mid \alpha_i \neq 0\}$ of Ω . Let D be a subcode of \mathbb{Z}_2^n . Set $D^{(0)} := \{\alpha \in D \mid \langle \alpha, \alpha \rangle = 0\}$ and $D^{(1)} := \{\alpha \in D \mid \langle \alpha, \alpha \rangle = 1\}$. For a code-word $\alpha = (\alpha_1, \dots, \alpha_n) \in D$, we define

$$U^\alpha := L(1/2, \alpha_1/2) \otimes \cdots \otimes L(1/2, \alpha_n/2).$$

Theorem 4.3. ([M2]) *For any linear code $D \subset \mathbb{Z}_2^n$, there exists a unique simple vertex operator superalgebra structure on $U_D := \bigoplus_{\alpha \in D} U^\alpha$ as an extension of $U^0 = L(1/2, 0)^{\otimes n}$. The even part is $U_{D^{(0)}} := \bigoplus_{\alpha \in D^{(0)}} U^\alpha$ and is a $D^{(0)}$ -graded simple current extension of U^0 , and the odd part is $U_{D^{(1)}} := \bigoplus_{\alpha \in D^{(1)}} U^\alpha$.*

Let D be an even subcode of \mathbb{Z}_2^n . The simple VOA U_D defined in the theorem above is called the *code VOA associated to a code D* . The representation theory of U_D is deeply studied in [M3]. We recall some results from [M3]. Since U_D is a D -graded simple current extension of a rational VOA $U^0 = L(1/2, 0)^{\otimes n}$, it is also rational. Let M be an irreducible U_D -module. Take an irreducible U^0 -submodule W of M . Then W is isomorphic to $\bigotimes_{i=1}^n L(1/2, h_i)$ with $h_i \in \{0, 1/2, 1/16\}$. Define the 1/16-word $\tau(W) = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_2^n$ of W by $\alpha_i = 1$ if and only if $h_i = 1/16$. Then by the fusion rules (3.1), $\tau(W)$ is determined independently of a choice of W and hence we can define the 1/16-word $\tau(M) \in \mathbb{Z}_2^n$ of M by $\tau(M) := \tau(W)$.

Assume that $\tau(M) = (0^n)$. Then W is isomorphic to $\bigotimes_{i=1}^n L(1/2, h_i)$ with $h_i \in \{0, 1/2\}$. We set $\gamma := (2h_1, \dots, 2h_n) \in \mathbb{Z}_2^n$. Since $L(1/2, 0)$ is a simple current $L(1/2, 0)$ -module, we have $D_W = 0$ and hence M is uniquely determined by W by Theorem 2.11 and has a shape

$$U_{D+\gamma} := \bigoplus_{\alpha \in D+\gamma} U^\alpha.$$

We call an irreducible U_D -module $U_{D+\gamma}$ a *coset module*. By Theorem 2.13 and the fusion rules (3.1), we have the following fusion rules for $\gamma_1, \gamma_2 \in \mathbb{Z}_2^n$:

$$U_{D+\gamma_1} \times U_{D+\gamma_2} = U_{D+\gamma_3}. \quad (4.2)$$

In particular, $U_{D+\delta} \times U_{D+\delta} = U_D$ for any $\delta \in \mathbb{Z}_2^n$ and hence all the coset modules are simple current U_D -modules by Lemma 2.6.

4.2 The Hamming code VOA

Let H_8 be the $[8, 4, 4]$ -Hamming code:

$$H_8 := \text{Span}_{\mathbb{Z}_2} \{(11111111), (11110000), (11001100), (10101010)\}.$$

It is well-known that H_8 is the unique doubly even self-dual linear code of length 8 up to isomorphism. Let us consider the Hamming code VOA U_{H_8} . In order to construct 2A-framed VOAs, we will need some special properties that the Hamming code VOA U_{H_8} possesses. Roughly speaking, we can identify $L(1/2, 1/16)$ with $L(1/2, 0)$ and $L(1/2, 1/2)$ by the symmetry of the Hamming code VOA.

Let X be an irreducible U_{H_8} -module whose top weight is in $\frac{1}{2}\mathbb{N}$. Then $\tau(X) = (0^8)$ or (1^8) and if $\tau(X) = (0^8)$ then $X \simeq U_{H_8+\gamma}$ for some $\gamma \in \mathbb{Z}_2^8$. If $\tau(X) = (1^8)$, then it is shown in [M3] that there is a unique linear character χ on H_8 such that

$$X \simeq \text{Ind}_{U_0}^{U_{H_8}}(L(1/2, 1/16)^{\otimes 8}, \chi) = L(1/2, 1/16)^{\otimes 8} \otimes_{\mathbb{C}} v_\chi, \quad (4.3)$$

where U^α , $\alpha \in H_8$, acts on $L(1/2, 1/16)^{\otimes 8}$ by the fusion rule (3.1) and on $\mathbb{C}v_\chi$ as a scalar $\chi(\alpha)$ according to the character χ . Since the dual group H_8^* of H_8 is naturally isomorphic to \mathbb{Z}_2^8/H_8 , we can find a unique coset $\delta_\chi + H_8 \in \mathbb{Z}_2^8/H_8$ such that $\chi(\alpha) = \langle \delta_\chi, \alpha \rangle$ for all $\alpha \in H_8$. So in the following we regard χ as an element in \mathbb{Z}_2^8 . Set $H(1/16, \chi) = L(1/2, 1/16)^{\otimes 8} \otimes_{\mathbb{C}} v_\chi$ for $\chi \in \mathbb{Z}_2^8$. Then $H(1/16, \chi_1) \simeq H(1/16, \chi_2)$ as U_{H_8} -modules if and only if $\chi_1 - \chi_2 \in H_8$ and the set of inequivalent irreducible U_{H_8} -modules whose top weights are contained in $\frac{1}{2}\mathbb{N}$ is given by

$$\{U_{H_8+\gamma}, H(1/16, \chi) \mid \gamma + H_8, \chi + H_8 \in \mathbb{Z}_2^8/H_8\}.$$

Surprisingly, we can identify a non-simple current $L(1/2, 0)^{\otimes 8}$ -module $L(1/2, 1/16)^{\otimes 8}$ with a coset module as follows:

Theorem 4.4. ([M4]) *For each $\chi \in \mathbb{Z}_2^8$, there is an automorphism $\sigma \in \text{Aut}(U_{H_8})$ such that the σ -conjugate module $H(1/16, \chi)^\sigma$ is isomorphic to a coset module $U_{H_8+\gamma}$ for some $\gamma \in \mathbb{Z}_2^8$ with $\langle \gamma, \gamma \rangle = 1$. In particular, $H(1/16, \chi)$ is a simple current U_{H_8} -module.*

Corollary 4.5. ([M4]) *As a \mathbb{Z}_2 -graded simple current extension of U_{H_8} , there is a unique simple SVOA structure on $U_{H_8} \oplus H(1/16, \chi)$ for all $\chi \in \mathbb{Z}_2^8$.*

Proof: We can take an irreducible coset U_{H_8} -module $U_{H_8+\gamma}$ with $\langle \gamma, \gamma \rangle = 1$ such that there is an automorphism $\sigma \in \text{Aut}(U_{H_8})$ such that the conjugate module $(U_{H_8+\gamma})^\sigma$ is isomorphic to $H(1/16, \chi)$ by Theorem 4.4. Then $U_{H_8} \oplus U_{H_8+\gamma}$ and $U_{H_8} \oplus H(1/16, \chi)$ form mutually conjugate \mathbb{Z}_2 -graded simple current extensions of U_{H_8} under $\sigma \in \text{Aut}(U_{H_8})$. Since $H_8 \cup (H_8 + \gamma)$ is an odd code, $U_{H_8} \oplus U_{H_8+\gamma}$ is a simple SVOA. Then so is $U_{H_8} \oplus H(1/16, \chi)$. ■

As an application of Proposition 4.4, the following fusion rules are established in [M4]:

Theorem 4.6. (*[M4]*) *We have the following fusion rules:*

$$\begin{aligned} U_{H_8+\alpha} \times U_{H_8+\beta} &= U_{H_8+\alpha+\beta}, \\ U_{H_8+\alpha} \times H(1/16, \beta) &= H(1/16, \alpha + \beta), \\ H(1/16, \alpha) \times H(1/16, \beta) &= U_{H_8+\alpha+\beta}, \end{aligned}$$

where $\alpha, \beta \in \mathbb{Z}_2^8$.

Thanks to Corollary 4.5 and Theorem 4.6, if an even linear code D contains many subcodes isomorphic to the Hamming code H_8 , then we can construct simple current extensions of the code VOA U_D by using Theorem 2.17.

4.3 Construction of 2A-framed VOA

In this subsection we give a refinement Miyamoto's construction of 2A-framed VOAs in [M4]. Here we assume the following:

Hypothesis 2.

- (1) (D, S) is a pair of even linear even codes of \mathbb{Z}_2^n such that
 - (1-i) $D \subset S^\perp$,
 - (1-ii) for each $\alpha \in S$, there is a subcode $E^\alpha \subset D$ such that E^α is a direct sum of the Hamming code H_8 and $\text{Supp}(E^\alpha) = \text{Supp}(\alpha)$, where $\text{Supp}(A)$ denotes $\cup_{\beta \in A} \text{Supp}(\beta)$ for a subset A of \mathbb{Z}_2^n .
- (2) $V^0 = U_D$ is the code VOA associated to the code D .
- (3) $\{V^\alpha \mid \alpha \in S\}$ is a set of irreducible V^0 -modules such that
 - (3-i) $\tau(V^\alpha) = \alpha$ for all $\alpha \in S$,
 - (3-ii) all V^α , $\alpha \in S$, have integral top weights,
 - (3-iii) the fusion product $V^\alpha \boxtimes_{V^0} V^\beta$ contains at least one $V^{\alpha+\beta}$. That is, there is a non-trivial V^0 -intertwining operator of type $V^\alpha \times V^\beta \rightarrow V^{\alpha+\beta}$ for any $\alpha, \beta \in S$.

Under Hypothesis 2 we will prove that $V := \oplus_{\alpha \in S} V^\alpha$ has a structure of an S -graded simple current extension of V^0 . Before we begin the proof, we prepare some lemmas.

Lemma 4.7. *Under Hypothesis 2, all V^α , $\alpha \in S$, are simple current V^0 -modules and we have the fusion rules $V^\alpha \times V^\beta = V^{\alpha+\beta}$ of V^0 -modules for all $\alpha, \beta \in S$.*

Proof: Suppose the fusion rule $V^\alpha \times V^\alpha = V^0$ of V^0 -modules holds. Then by Lemma 2.6, V^α is a simple current V^0 -module because $V^0 = U_D$ is a rational C_2 -cofinite VOA of CFT-type. Then by Hypothesis 2 (3-iii) we have the desired fusion rule $V^\alpha \times V^\beta = V^{\alpha+\beta}$. Therefore, we only prove the fusion rule $V^\alpha \times V^\alpha = V^0$ for each $\alpha \in S$. By Hypothesis 2

(1-i), D contains a subcode E^α which is isomorphic to a direct sum of H_8 and $\text{Supp}(E^\alpha) = \text{Supp}(\alpha)$. We may assume that $\alpha = (1^{8s}0^t)$ with $8s + t = n$. Then U_D contains a sub VOA

$$L := U_{E^\alpha} \otimes L(1/2, 0)^{\otimes t} \simeq (U_{H_8})^{\otimes s} \otimes L(1/2, 0)^{\otimes t}$$

and V^α contains an irreducible L -submodule X isomorphic to

$$H(1/16, \chi_1) \otimes \cdots H(1/16, \chi_s) \otimes L(1/2, h_1) \otimes \cdots \otimes L(1/2, h_t)$$

with $\chi_i \in \mathbb{Z}_2^8$, $1 \leq i \leq s$, and $h_j \in \{0, 1/2\}$, $1 \leq j \leq t$. Let $D = \sqcup_{i=0}^k (E^\alpha + \beta_i)$ be a coset decomposition. We write $\beta_i = \gamma_i + \delta_i$ such that $\text{Supp}(\gamma_i) \subset \text{Supp}(\alpha)$ and $\text{Supp}(\delta_i) \cap \text{Supp}(\alpha) = \emptyset$. Then $U_{E^\alpha + \beta_i}$ is isomorphic to

$$U_{E^\alpha + \gamma_i} \otimes L(1/2, (\delta_i)_{8s+1}/2) \otimes \cdots \otimes L(1/2, (\delta_i)_n/2)$$

as an L -module and $U_D = \oplus_{i=1}^k U_{E^\alpha + \beta_i}$ is a D/E^α -graded simple current extension of L . Then by the fusion rules (3.1) and Theorem 4.6, $(U_{E^\alpha + \beta_i}) \boxtimes_L X$ is an irreducible L -module and $(U_{E^\alpha + \beta_i}) \boxtimes_L X \not\simeq (U_{E^\alpha + \beta_j}) \boxtimes_L X$ unless $i = j$. Therefore, V^α as an L -module is isomorphic to $V^\alpha = \oplus_{i=1}^k (U_{E^\alpha + \beta_i}) \boxtimes_L X$. Namely, V^α is a D/E^α -stable U_D -module. Then by Theorem 2.13 together with fusion rules (3.1) and Theorem 4.6 we have a fusion rule $V^\alpha \times V^\alpha = V^0$ of U_D -modules which is a lifting of the fusion rule $X \times X = L$ of L -modules. \blacksquare

Lemma 4.8. *Under Hypothesis 2, the space $V^0 \oplus V^\alpha$ forms a simple VOA as a \mathbb{Z}_2 -graded simple current extension of V^0 for each $\alpha \in S \setminus 0$.*

Proof: Here we use the same notation as in the proof of Lemma 4.7. By the coset decomposition $D = \sqcup_{i=1}^k (E^\alpha + \beta_i)$, $V^0 = U_D = \oplus_{i=1}^k U_{E^\alpha + \beta_i}$ is a D/E^α -graded simple current extension of $L = U_{E^\alpha} \otimes L(1/2, 0)^{\otimes t} \simeq (U_{H_8})^{\otimes s} \otimes L(1/2, 0)^{\otimes t}$. By the fusion rule $X \times X = L$ of L -modules, X is a simple current L -module by Lemma 2.6. Then by the associativity of fusion products (cf. [H4]), $(U_{E^\alpha + \beta_i}) \boxtimes_L X$ is also a simple current L -module. Thus we obtain the set of inequivalent simple current L -modules

$$\mathcal{S} = \{ U_{E^\alpha + \beta_i}, (U_{E^\alpha + \beta_j}) \boxtimes_L X \mid 1 \leq i, j \leq k \}$$

with the following $((D/E^\alpha) \oplus \mathbb{Z}_2)$ -graded fusion rules:

$$\begin{aligned} U_{E^\alpha + \beta_i} \times U_{E^\alpha + \beta_j} &= U_{E^\alpha + \beta_i + \beta_j}, \\ U_{E^\alpha + \beta_i} \times (U_{E^\alpha + \beta_j} \boxtimes_L X) &= (U_{E^\alpha + \beta_i + \beta_j}) \boxtimes_L X, \\ (U_{E^\alpha + \beta_i} \boxtimes_L X) \times (U_{E^\alpha + \beta_j} \boxtimes_L X) &= U_{E^\alpha + \beta_i + \beta_j}. \end{aligned}$$

Since $U_D = \oplus_{i=1}^k U_{E^\alpha + \beta_i}$ has a structure of a D/E^α -graded simple current extension of L and $L \oplus X$ has a structure of a \mathbb{Z}_2 -graded simple current extension of L by Corollary 4.5,

we can apply Theorem 2.17 to \mathcal{S} and hence we obtain a $((D/E^\alpha) \oplus \mathbb{Z}_2)$ -graded simple current extension

$$\left\{ \bigoplus_{i=1}^k U_{E^\alpha + \beta_i} \right\} \bigoplus \left\{ \bigoplus_{i=1}^k (U_{E^\alpha + \beta_i}) \boxtimes_L X \right\}$$

of L . Since $V^0 = \bigoplus_{i=1}^k U_{E^\alpha + \beta_i}$ and $V^\alpha = \bigoplus_{i=1}^k (U_{E^\alpha + \beta_i}) \boxtimes_L X$, the \mathbb{Z}_2 -graded space $V^0 \oplus V^\alpha$ carries a simple VOA structure which is the desired \mathbb{Z}_2 -graded simple current extension of V^0 . \blacksquare

Now we can prove

Theorem 4.9. ([M4]) *Under Hypothesis 2, the space $V = \bigoplus_{\alpha \in S} V^\alpha$ has a unique structure of a simple VOA as an S -graded simple current extension of V^0 . In particular, there exists a 2A-framed VOA whose structure codes are (D, S) .*

Proof: Let $\{\alpha_1, \dots, \alpha_r\}$ be a linear basis of S and set $S^i := \text{Span}_{\mathbb{Z}_2}\{\alpha_1, \dots, \alpha_i\}$ for $1 \leq i \leq r$. We proceed by induction on r . The case $r = 0$ is trivial and the case $r = 1$ is given by Lemma 4.8. Now assume that $\bigoplus_{\beta \in S^i} V^\beta$ has a structure of a simple VOA for $1 \leq i \leq r - 1$. Then the set

$$\mathcal{T} = \{ V^\beta, V^{\beta + \alpha_{i+1}} \mid \beta \in S^i \}$$

consists of inequivalent simple current V^0 -modules with $(S^i \oplus \mathbb{Z}_2) = S^{i+1}$ -graded fusion rules:

$$V^{\beta_1} \times V^{\beta_2} = V^{\beta_1 + \beta_2}, \quad V^{\beta_1} \times V^{\beta_2 + \alpha_{i+1}} = V^{\beta_1 + \beta_2 + \alpha_{i+1}}, \quad V^{\beta_1 + \alpha_{i+1}} \times V^{\beta_2 + \alpha_{i+1}} = V^{\beta_1 + \beta_2},$$

where $\beta_1, \beta_2 \in S^i$. By inductive assumption, $\bigoplus_{\beta \in S^i} V^\beta$ is an S^i -graded simple current extension of V^0 , and by Lemma 4.8, a direct sum $V^0 \oplus V^{\alpha_{i+1}}$ becomes a \mathbb{Z}_2 -graded simple current extension of V^0 . Therefore, we can apply Theorem 2.17 to \mathcal{T} to obtain the S^{i+1} -graded simple current extension $\bigoplus_{\beta \in S^{i+1}} V^\beta$ of V^0 . Repeating this procedure, we finally obtain $S^r = S$ -graded simple current extension $V = \bigoplus_{\alpha \in S} V^\alpha$ of $V^0 = U_D$. \blacksquare

Remark 4.10. In [M4], Miyamoto assumed stronger conditions than that in Hypothesis 2. In particular, he assumed that the structure codes (D, S) are of length $8k$ for some positive integer k . Our refinement enable us to construct 2A-framed VOAs with structure codes of any length as long as Hypothesis 2 is satisfied.

Extension to SVOA. Let V be a 2A-framed VOA with structure codes (D, S) . Then by definition we have a decomposition $V = \bigoplus_{\alpha \in S} V^\alpha$ such that $V^0 \simeq U_D$ and $\tau(V^\alpha) = \alpha$. Assume that the pair (D, S) satisfies the condition (1) of Hypothesis 2. Then V is an S -graded simple current extension of V^0 by Lemma 4.7. Suppose that there is a vector $\gamma \in S^\perp \setminus D$ such that $\langle \gamma, \gamma \rangle = 1$. Since the powers of z in an $L(1/2, 0)$ -intertwining

operator of type $L(1/2, 1/2) \times L(1/2, 1/16) \rightarrow L(1/2, 1/16)$ are half-integral, the powers of z in a U_D -intertwining operator of type $U_{D+\gamma} \times V^\alpha \rightarrow U_{D+\gamma} \boxtimes_{U_D} V^\alpha$ are integral for all $\alpha \in S$. Therefore, all $V^\alpha \boxtimes_{U_D} U_{D+\gamma}$, $\alpha \in S$, have half-integral top weights. Since the $1/16$ -word of $V^\alpha \boxtimes_{U_D} U_{D+\gamma}$ is α by the fusion rules (3.1), the induced module

$$\text{Ind}_{V^0}^V U_{D+\gamma} = \bigoplus_{\alpha \in S} V^\alpha \boxtimes_{V^0} U_{D+\gamma}$$

has a unique V -module structure by Theorem 2.11. Since $U_D \oplus U_{D+\gamma}$ is a simple current super-extension of $V^0 = U_D$ by Theorem 4.3, by applying Theorem 2.17 we obtain:

Theorem 4.11. *With reference to the setup above, $V \oplus \text{Ind}_{V^0}^V U_{D+\gamma}$ forms a simple current super-extension of V .*

5 The baby-monster SVOA

As shown in [DMZ], the moonshine VOA V^\natural has a 2A-frame $\omega^\natural = e^1 + \dots + e^{48}$. One of its structure codes are determined in [DGH] and [M5]. Let S be the Reed-Müller code $RM(4, 1)$ defined as follows:

$$S = \text{Span}_{\mathbb{Z}_2} \{(1^{16}), (1^8 0^8), (1^4 0^4 1^4 0^4), (1^2 0^2 1^2 0^2 1^2 0^2), (1010101010101010)\} \subset \mathbb{Z}_2^{16}.$$

Then define

$$S^\natural := \{(\alpha, \alpha, \alpha), (\alpha^c, \alpha, \alpha), (\alpha, \alpha^c, \alpha), (\alpha, \alpha, \alpha^c) \in \mathbb{Z}_2^{48} \mid \alpha \in S, \alpha^c := \alpha + (1^{16})\}$$

and $D^\natural := (S^\natural)^\perp$.

Theorem 5.1. *([DGH] [M5]) The moonshine VOA V^\natural has a structure codes (D^\natural, S^\natural) .*

One can easily check that the pair (D^\natural, S^\natural) satisfies the condition (1) of Hypothesis 2. Thus, by Lemma 4.7, we have

Corollary 5.2. *Let $V^\natural = \bigoplus_{\alpha \in S^\natural} (V^\natural)^\alpha$ be the $1/16$ -word decomposition according to the structure codes (D^\natural, S^\natural) . Then the pair (D^\natural, S^\natural) and the set $\{(V^\natural)^\alpha \mid \alpha \in S^\natural\}$ satisfy Hypothesis 2. Therefore, V^\natural is an S^\natural -graded simple current extension of $(V^\natural)^0 = U_{D^\natural}$.*

Now set $e = e^1$ and consider the commutant subalgebra $T_e^\natural(0)$ of $\text{Vir}(e)$ in V^\natural . Since $\{1\} \cap \text{Supp}(S^\natural) \neq \emptyset$, $V_e^\natural(1/16)$ is not zero. Then by the condition (1) of Hypothesis 2, $V_e(1/2)$ is not zero, too. Therefore, we obtain a decomposition

$$V^\natural = L(1/2, 0) \otimes T_e^\natural(0) \oplus L(1/2, 1/2) \otimes T_e^\natural(1/2) \oplus L(1/2, 1/16) \otimes T_e^\natural(1/16)$$

such that $T_e^{\natural}(h) \neq 0$ for $h = 0, 1/2, 1/16$. By Theorem 3.6, we know that $T_e^{\natural}(0) \oplus T_e^{\natural}(1/2)$ has a structure of a simple vertex operator superalgebra. This algebra was first considered by Höhn [H1] and he called it the *baby-monster SVOA*, because the centralizer $C_{\text{Aut}(V^{\natural})}(\tau_e)$ is isomorphic to the 2-fold central extension $\langle \tau_e \rangle \cdot \mathbb{B}$ of the baby-monster sporadic finite simple group \mathbb{B} [ATLAS] and so \mathbb{B} naturally acts on it. Following him, we set $VB^0 := T_e^{\natural}(0)$, $VB^1 := T_e^{\natural}(1/2)$ and $VB := T_e^{\natural}(0) \oplus T_e^{\natural}(1/2)$. We also know that $T_e^{\natural}(1/16)$ is an irreducible \mathbb{Z}_2 -twisted VB -module by Theorem 3.8, and so we set $VB_T := T_e^{\natural}(1/16)$ for convention. Since all the conformal vectors of V^{\natural} with central charge $1/2$ are conjugate under the Monster $\mathbb{M} = \text{Aut}(V^{\natural})$ by [C] and [M1], the algebraic structures on VB and VB_T are independent of choice a conformal vector $e = e^1$.

By definition, the Virasoro vector of VB^0 is given by $\omega^{\natural} - e^1 = e^2 + \cdots + e^{48}$. Thus VB^0 is a 2A-framed VOA. We compute the structure codes of VB^0 . Set

$$(S^{\natural})^0 := \{\alpha \in S^{\natural} \mid \{1\} \cap \text{Supp}(\alpha) = \emptyset\}, \quad (S^{\natural})^1 := \{\alpha \in S^{\natural} \mid \{1\} \cap \text{Supp}(\alpha) = \{1\}\}.$$

Then $S^{\natural} = (S^{\natural})^0 \sqcup (S^{\natural})^1$ and by definition of Miyamoto involution, $(V^{\natural})^{\langle \tau_e \rangle} = \bigoplus_{\alpha \in (S^{\natural})^0} (V^{\natural})^{\alpha}$ and $V_e^{\natural}(1/16) = \bigoplus_{\beta \in (S^{\natural})^1} (V^{\natural})^{\beta}$. Since V^{\natural} is an S^{\natural} -graded simple current extension of $(V^{\natural})^0$, the fixed point subalgebra $(V^{\natural})^{\langle \tau_e \rangle}$ is also an $(S^{\natural})^0$ -graded simple current extension of $(V^{\natural})^0$ and $V_e^{\natural}(1/16)$ is a simple current $(V^{\natural})^{\langle \tau_e \rangle}$ -module by Corollary 2.14. Now define $\phi_{\epsilon} : \mathbb{Z}_2^{47} \hookrightarrow \mathbb{Z}_2^{48}$ by $\mathbb{Z}_2^{47} \ni \alpha \mapsto (\epsilon, \alpha) \in \mathbb{Z}_2^{48}$ for $\epsilon = 0, 1$, and set

$$D^{b,\epsilon} := \{\alpha \in \mathbb{Z}_2^{47} \mid \phi_{\epsilon}(\alpha) \in D^{\natural}\}, \quad \epsilon = 0, 1, \quad S^b := \{\beta \in \mathbb{Z}_2^{47} \mid \phi_0(\beta) \in (S^{\natural})^0\}.$$

Proposition 5.3. *The structure codes of VB^0 with respect to the 2A-frame $e^2 + \cdots + e^{48}$ are $(D^{b,0}, S^b)$.*

Proof: For $\alpha \in (S^{\natural})^0$, define $(V^{\natural})^{\alpha,\epsilon}$ to be the sum of all irreducible $\bigotimes_{i=1}^{48} \text{Vir}(e^i)$ -submodules of $(V^{\natural})^{\alpha}$ whose $\text{Vir}(e^1)$ -components are isomorphic to $L(1/2, \epsilon/2)$ for $\epsilon = 0, 1$. Then $V_e^{\natural}(1/2) \neq 0$ implies that $(V^{\natural})^{\alpha,\epsilon} \neq 0$ for all $\alpha \in (S^{\natural})^0$ and $\epsilon = 0, 1$. Therefore, $(V^{\natural})^{\alpha} = (V^{\natural})^{\alpha,0} \oplus (V^{\natural})^{\alpha,1}$ and we obtain 1/16-word decompositions $V_e^{\natural}(0) = \bigoplus_{\alpha \in (S^{\natural})^0} (V^{\natural})^{\alpha,0}$ and $V_e^{\natural}(1/2) = \bigoplus_{\alpha \in (S^{\natural})^0} (V^{\natural})^{\alpha,1}$. Since $D^{\natural} = \phi_0(D^{b,0}) \sqcup \phi_1(D^{b,1})$, $(V^{\natural})^{0,0} \simeq L(1/2, 0) \otimes U_{D^{b,0}}$. Thus VB^0 has a 1/16-word decomposition $VB^0 = \bigoplus_{\alpha \in S^b} (VB^0)^{\alpha}$ such that $\tau((VB^0)^{\alpha}) = \alpha$ and $(VB^0)^0 \simeq U_{D^{b,0}}$. Hence the structure codes of VB^0 are $(D^{b,0}, S^b)$. \blacksquare

Remark 5.4. By the proof above, we find that VB^1 also has a 1/16-word decomposition $VB^1 = \bigoplus_{\alpha \in S^b} (VB^1)^{\alpha}$ such that $\tau((VB^1)^{\alpha}) = \alpha$. In particular, $(VB^1)^0$ is isomorphic to a coset module $U_{D^{b,1}}$.

The following is easy to see:

Lemma 5.5. *The pair $(D^{b,0}, S^b)$ satisfies the condition (1) of Hypothesis 2.*

Therefore, $VB^0 = \oplus_{\alpha \in S^b} (VB^0)^\alpha$ is an S^b -graded simple current extension of the code VOA $U_{D^b,0}$. Since the pair (D^b, S^b) and the set $\{(VB^0)^\alpha \mid \alpha \in S^b\}$ satisfy Hypothesis 2, we can construct VB^0 without reference to V^\natural by Theorem 4.9.

Proposition 5.6. *VB^1 is a simple current VB^0 -module.*

Proof: By Remark 5.4, VB^1 is isomorphic to the induced module $\text{Ind}_{U_{D^b,0}}^{VB^0} U_{D^b,1}$ which is an S^b -stable VB^0 -module. Therefore, we have the fusion rule

$$VB^1 \times VB^1 = \text{Ind}_{U_{D^b,0}}^{VB^0} (U_{D^b,1} \boxtimes_{U_{D^b,0}} U_{D^b,1}) = \text{Ind}_{U_{D^b,0}}^{VB^0} U_{D^b,0} = VB^0$$

by Theorem 2.13. Thus VB^1 is a simple current VB^0 -module by Lemma 2.6. ■

Remark 5.7. We note that by using Theorem 4.11 we can define the SVOA structure on VB without reference to V^\natural .

Up to now, we have established that V^\natural and its conformal vector e satisfy all the conditions in Hypothesis 1. Moreover, it is shown in [C] and [M1] that there is a one-to-one correspondence between the set of conformal vectors of V^\natural with central charge $1/2$ and the set of corresponding Miyamoto involutions on V^\natural which is known to be the 2A-conjugacy class of the Monster. Therefore, $C_{\text{Aut}(V^\natural)}(e) = C_{\mathbb{M}}(\tau_e) \simeq \langle \tau_e \rangle \cdot \mathbb{B}$. Thus, $C_{\text{Aut}(V^\natural)}(e)/\langle \tau_e \rangle$ is a simple group and we can apply Theorem 3.14 to VB .

Theorem 5.8. (1) *The SVOA VB obtained from V^\natural by cutting off the Ising model is a simple SVOA.*

(2) *The piece VB_T obtained from V^\natural is an irreducible \mathbb{Z}_2 -twisted VB -module.*

(3) *$\text{Aut}(VB^0) \simeq \mathbb{B}$ and $\text{Aut}(VB) \simeq 2 \times \mathbb{B}$.*

(4) *VB_T as a VB^0 -module is irreducible. Thus, there are exactly three irreducible VB^0 -modules, VB^0 , VB^1 and VB_T .*

(5) *The fusion rules for irreducible VB^0 -modules are as follows:*

$$VB^1 \times VB^1 = VB^0, \quad VB^1 \times VB_T = VB_T, \quad VB_T \times VB_T = VB^0 + VB^1.$$

Proof: (1) follows from Theorem 3.6, (2) follows from 3.8 and (3) will follow from Theorem 3.14 and the fact $C_{\text{Aut}(V^\natural)}(e) = \langle \tau_e \rangle \cdot \mathbb{B}$.

Consider (4). By Corollary 3.15, VB_T as a VB^0 -module is irreducible. Then the assertion follows from Proposition 3.11.

Consider (5). We only have to show the fusion rule $VB_T \times VB_T = VB^0 + VB^1$. By considering the $1/16$ -word of VB_T , the fusion product $VB_T \times VB_T$ is contained in $\text{NVB}^0 \oplus \text{NVB}^1$ in the fusion algebra for VB^0 . Write $VB_T \times VB_T = n_0 VB^0 + n_1 VB^1$ with $n_0, n_1 \in \mathbb{N}$. Then the simplicity of V^\natural implies $n_0 \neq 0$ and $n_1 \neq 0$. And by applying VB^1 to $VB_T \times VB_T$, we see that $n_0 = n_1$. Since the dual module of VB_T is isomorphic to VB_T , the space of VB^0 -intertwining operator of type $VB_T \times VB_T \rightarrow VB^0$ is one-dimensional. Thus $n_0 = n_1 = 1$ as desired. ■

Remark 5.9. The assertion (1) of Theorem 5.8 is already shown by Höhn in [Hö1], and (3) of Theorem 5.8 is also proved in [Hö2]. However, Höhn's proofs in [Hö1] [Hö2] and ours are quite different. In particular, in [Hö2], he used many results on the baby-monster simple group. Our argument can be applied to any 2A-framed VOAs satisfying Hypothesis 1 and Hypothesis 2 since we have only used the facts that $C_{\text{Aut}(V^\natural)}(e) = C_{\text{Aut}(V^\natural)}(\tau_e)$ and $C_{\text{Aut}(V^\natural)}(\tau_e)/\langle \tau_e \rangle$ is a simple group.

The classification of irreducible VB^0 -modules has many interesting corollaries.

Corollary 5.10. *The irreducible 2A-twisted V^\natural -module has a shape*

$$L(1/2, 1/2) \otimes VB^0 \oplus L(1/2, 0) \otimes VB^1 \oplus L(1/2, 1/16) \otimes VB_T.$$

Proof: Follows from Theorem 5.8, Theorem 3.10 and Proposition 3.11. ■

Remark 5.11. A straightforward construction of the 2A-twisted (and 2B-twisted) V^\natural -module is given by Lam [L2]. In his construction, it is given as $U_{D^\natural+\gamma} \boxtimes_{U_{D^\natural}} V^\natural$ with $\gamma = (10^{47}) \in (\mathbb{Z}/2\mathbb{Z})^{48}$.

Corollary 5.12. *For any conformal vector $e \in V^\natural$ with central charge $1/2$, there is no automorphism ρ on V^\natural such that $\rho(V_e^\natural(h)) = V_e^\natural(h)$ for $h = 0, 1/2$ and $\rho|_{(V^\natural)^{\langle \tau_e \rangle}} = \sigma_e$.*

Proof: Suppose such an automorphism ρ exists. We remark that ρ also preserves the space $V_e(1/16)$ as $\rho \in C_{\text{Aut}(V)}(e)$. We view $V_e^\natural(1/16)$ as a $(V^\natural)^{\langle \tau_e \rangle}$ -module by a restriction of the vertex operator map $Y_{V^\natural}(\cdot, z)$ on V^\natural . Consider the σ_e -conjugate $(V^\natural)^{\langle \tau_e \rangle}$ -module $V_e^\natural(1/16)^{\sigma_e}$. By Theorem 5.8 and Proposition 3.11, $V_e^\natural(1/16)^{\sigma_e}$ is not isomorphic to $V_e^\natural(1/16)$ as a $(V^\natural)^{\langle \tau_e \rangle}$ -module. On the other hand, we can take a canonical linear isomorphism $\varphi : V_e^\natural(1/16) \rightarrow V_e^\natural(1/16)^{\sigma_e}$ such that $Y_{V_e^\natural(1/16)^{\sigma_e}}(a, z)\varphi v = \varphi Y_{V^\natural}(\sigma_e a, z)v$ for all $a \in (V^\natural)^{\langle \tau_e \rangle}$ and $v \in V_e^\natural(1/16)$ by definition of the conjugate module. Then we have

$$Y_{V_e^\natural(1/16)^{\sigma_e}}(a, z)\varphi \rho v = \varphi Y_{V^\natural}(\sigma_e a, z)\rho v = \varphi Y_{V^\natural}(\rho a, z)\rho v = \varphi \rho Y_{V^\natural}(a, z)v$$

for any $a \in (V^\natural)^{\langle \tau_e \rangle}$ and $v \in V_e^\natural(1/16)$. Thus $\varphi \rho$ defines a $(V^\natural)^{\langle \tau_e \rangle}$ -isomorphism between $V_e^\natural(1/16)$ and $V_e^\natural(1/16)^{\sigma_e}$, which is a contradiction. ■

Corollary 5.13. *The 2A-orbifold construction applied to the moonshine VOA V^\natural yields V^\natural itself again.*

Proof: Follows from Theorem 5.8 and Corollary 3.15. ■

Remark 5.14. The statement in the corollary above was conjectured by Tuite [Tu]. In [Tu], Tuite has shown that any \mathbb{Z}_p -orbifold construction of V^\natural yields the moonshine VOA V^\natural or the Leech lattice VOA V_Λ under the uniqueness conjecture of the moonshine VOA which states that V^\natural constructed by Frenkel et. al. [FLM] is the unique holomorphic VOA with central charge 24 whose weight one subspace is trivial.

Finally, we end this paper by presenting the modular transformations of characters of VB^0 -modules. Here the character means the conformal character, not the q -dimension, of modules. Recall the characters of $L(1/2, 0)$ -modules. By our explicit construction in Section 3.1, one can easily prove the following (cf. [FFR] [FRW]):

$$\begin{aligned}\mathrm{ch}_{L(1/2,0)}(\tau) &= (1/2) \cdot q^{-1/48} \left\{ \prod_{n=0}^{\infty} (1 + q^{n+1/2}) + \prod_{n=0}^{\infty} (1 - q^{n+1/2}) \right\}, \\ \mathrm{ch}_{L(1/2,1/2)}(\tau) &= (1/2) \cdot q^{-1/48} \left\{ \prod_{n=0}^{\infty} (1 + q^{n+1/2}) - \prod_{n=0}^{\infty} (1 - q^{n+1/2}) \right\}, \\ \mathrm{ch}_{L(1/2,1/16)}(\tau) &= q^{-1/24} \prod_{n=1}^{\infty} (1 + q^n).\end{aligned}$$

The following modular transformations are well-known:

$$\begin{aligned}\mathrm{ch}_{L(1/2,0)}(-1/\tau) &= \frac{1}{2} \mathrm{ch}_{L(1/2,0)}(\tau) + \frac{1}{2} \mathrm{ch}_{L(1/2,1/2)}(\tau) + \frac{1}{\sqrt{2}} \mathrm{ch}_{L(1/2,1/16)}(\tau), \\ \mathrm{ch}_{L(1/2,1/2)}(-1/\tau) &= \frac{1}{2} \mathrm{ch}_{L(1/2,0)}(\tau) + \frac{1}{2} \mathrm{ch}_{L(1/2,1/2)}(\tau) - \frac{1}{\sqrt{2}} \mathrm{ch}_{L(1/2,1/16)}(\tau), \\ \mathrm{ch}_{L(1/2,1/16)}(-1/\tau) &= \frac{1}{\sqrt{2}} \mathrm{ch}_{L(1/2,0)}(\tau) - \frac{1}{\sqrt{2}} \mathrm{ch}_{L(1/2,1/2)}(\tau).\end{aligned}$$

Set $j(\tau) := J(\tau) - 744$, where $J(\tau)$ is the famous $\mathrm{SL}_2(\mathbb{Z})$ -invariant. Since $\mathrm{ch}_{V^\natural}(\tau) = j(\tau)$ and

$$\mathrm{ch}_{V^\natural}(\tau) = \mathrm{ch}_{L(1/2,0)}(\tau) \mathrm{ch}_{VB^0}(\tau) + \mathrm{ch}_{L(1/2,1/2)}(\tau) \mathrm{ch}_{VB^1}(\tau) + \mathrm{ch}_{L(1/2,1/16)}(\tau) \mathrm{ch}_{VB_T}(\tau),$$

we can write down the characters of irreducible VB^0 -modules by using those of V^\natural and $L(1/2, 0)$ -modules. This computation is already done in [Ma] by using Norton's trace formula. The results are written as a rational expression involving the functions $j(\tau)$, $\mathrm{ch}_{L(1/2,h)}(\tau)$, $h = 0, 1/2, 1/16$, their first and second derivatives and the Eisenstein series $E_2(\tau)$ and $E_4(\tau)$, see [Ma].

By Zhu's theorem [Z], the linear space spanned by $\{\mathrm{ch}_{VB^0}(\tau), \mathrm{ch}_{VB^1}(\tau), \mathrm{ch}_{VB_T}(\tau)\}$ affords an $\mathrm{SL}_2(\mathbb{Z})$ -action. By using the modular transformations for $j(\tau)$ and $\mathrm{ch}_{L(1/2,h)}(\tau)$, $h = 0, 1/2, 1/16$, we can show the following modular transformations:

$$\begin{aligned}\mathrm{ch}_{VB^0}(-1/\tau) &= \frac{1}{2} \mathrm{ch}_{VB^0}(\tau) + \frac{1}{2} \mathrm{ch}_{VB^1}(\tau) + \frac{1}{\sqrt{2}} \mathrm{ch}_{VB_T}(\tau), \\ \mathrm{ch}_{VB^1}(-1/\tau) &= \frac{1}{2} \mathrm{ch}_{VB^0}(\tau) + \frac{1}{2} \mathrm{ch}_{VB^1}(\tau) - \frac{1}{\sqrt{2}} \mathrm{ch}_{VB_T}(\tau), \\ \mathrm{ch}_{VB_T}(-1/\tau) &= \frac{1}{\sqrt{2}} \mathrm{ch}_{VB^0}(\tau) - \frac{1}{\sqrt{2}} \mathrm{ch}_{VB^1}(\tau).\end{aligned}$$

Namely, we have exactly the same modular transformation laws for the Ising model $L(1/2, 0)$. As in Theorem 5.8, we also note that the fusion algebra for VB^0 is also canonically isomorphic to that of $L(1/2, 0)$. Therefore, we may say that $L(1/2, 0)$ and VB^0 form a dual-pair in the moonshine VOA V^\natural .

Acknowledgments

The author wishes to thank the members of the Komaba Seminar on Finite Groups for valuable discussions and for noticing a gap in a prototype version of this paper. He also thank Professor Atsushi Matsuo for the information of Tuite's work and Professor Ching Hung Lam for his suggestions improving this paper.

References

- [ATLAS] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, ATLAS of finite groups, Clarendon Press, Oxford, 1985.
- [C] J. H. Conway, A simple construction for the Fischer-Griess monster group, *Invent. Math.* **79** (1985), 513–540.
- [DGH] C. Dong, R.L. Griess and G. Höhn, Framed vertex operator algebras, codes and the moonshine module, *Comm. Math. Phys.* **193** (1998), 407–448.
- [DL] C. Dong and J. Lepowsky, Generalized vertex algebras and relative vertex operators, Progress in Math. **112**, Birkhäuser, Boston, 1993.
- [DLM1] C. Dong, H. Li and G. Mason, Modular-invariance of trace functions in orbifold theory and generalized moonshine, *Comm. Math. Phys.* **214** (2000), 1–56.
- [DLM2] C. Dong, H. Li and G. Mason, Hom functor and the associativity of tensor products of modules for Vertex operator algebras, *J. Algebra* **188** (1997), 443–475.
- [DLM3] C. Dong, H. Li and G. Mason, Simple currents and extensions of vertex operator algebras, *Comm. Math. Phys.* **180**, 671–707.
- [DM1] C. Dong and G. Mason, On quantum Galois theory, *Duke Math. J.* **86** (1997), 305–321.
- [DM2] C. Dong and G. Mason, Rational vertex operator algebras and the effective central charge, math.QA/0201318.
- [DMZ] C. Dong, G. Mason and Y. Zhu, Discrete series of the Virasoro algebra and the moonshine module, Proc. Symp. Pure. Math., American Math. Soc. **56** II (1994), 295–316.
- [FFR] A. J. Feingold, I. B. Frenkel and J. F.X. Ries, Spinor construction of vertex operator algebras, triality, and $E_8^{(1)}$, *Contemp. Math.* **121** (1991).
- [FLM] I.B. Frenkel, J. Lepowsky and A. Meurman, Vertex Operator Algebras and the Monster, Academic Press, New York, 1988.
- [FRW] A. J. Feingold, J. F.X. Ries and M. Weiner, Spinor construction of the $c = \frac{1}{2}$ minimal model. Moonshine, the Monster, and related topics (South Hadley, MA, 1994), 45–92, Contemp. Math., 193, Amer. Math. Soc., Providence, RI, 1996.

- [FZ] I.B. Frenkel and Y. Zhu, Vertex operator algebras associated to representation of affine and Virasoro algebras, *Duke Math. J.* **66** (1992), 123–168.
- [G] R.L. Griess, The friendly giant, *Invent. Math.* **69** (1982), 1–102.
- [Hö1] G. Höhn, Selbstduale Vertexoperatorsuperalgebren und das Babymonster, Ph.D. thesis, Bonn 1995.
- [Hö2] G. Höhn, The group of symmetries of the shorter moonshine module, [math.QA/0210076](#).
- [H1] Y.-Z. Huang, A theory of tensor products for module categories for a vertex operator algebra, IV, *J. Pure Appl. Algebra* **100** (1995), 173–216.
- [H2] Y.-Z. Huang, Virasoro vertex operator algebras, (non-meromorphic) operator product expansion and the tensor product theory, *J. Algebra* **182** (1996), 201–234.
- [H3] Y.-Z. Huang, A nonmeromorphic extension of the moonshine module vertex operator algebra, in "Moonshine, the Monster and Related Topics, Proc. Joint Summer Research Conference, Mount Holyoke, 1994" (C. Dong and G. Mason, Eds.), *Contemporary Math.*, pp. 123–148, Amer. Math. Soc., Providence, RI, 1996.
- [H4] Y.-Z. Huang, Differential equations and intertwining operators, [math.QA/0206206](#).
- [HL1] Y.-Z. Huang and J. Lepowsky, Toward a theory of tensor product for representations of a vertex operator algebra, in Proc. 20th Intl. Conference on Diff. Geom. Methods in Theoretical Physics, New York, 1991, ed. S. Catto and A. Rocha, World Scientific, Singapore, 1992, Vol. 1, 344–354.
- [HL2] Y.-Z. Huang and J. Lepowsky, Tensor products of modules for a vertex operator algebra and vertex tensor categories, in: *Lie Theory and Geometry*, ed. R. Brylinski, J.-L. Brylinski, V. Guillemin, V. Kac, Birkhauser, Boston, 1994, 349–383.
- [HL3] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for module categories for a vertex operator algebra, I, *Sel. Math.* **1** (1995), 699–756.
- [HL4] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for module categories for a vertex operator algebra, III, *J. Pure Appl. Alg.* **100** (1995), 141–171.
- [KR] V. G. Kac and A. K. Raina, "Bombay Lectures on Highest Weight Representations of Infinite Dimensional Lie algebras," World Scientific, Singapore, 1987.
- [L1] C. H. Lam, Induced modules for orbifold vertex operator algebras, *J. Math. Soc. Japan* **53** (2001), 541–557.
- [L2] C. H. Lam, Some twisted modules for framed vertex operator algebras, *J. Algebra* **231** (2000), 331–341.
- [LYY] C. H. Lam, H. Yamada and H. Yamauchi, Vertex operator algebras, extended E_8 -diagram, and McKay's observation on the Monster simple group, preprint.

- [Li1] H. Li, Local systems of vertex operators, vertex superalgebras and modules, *J. Pure Appl. Algebra* **109** (1996), 143–195.
- [Li2] H. Li, Local systems of twisted vertex operators, vertex operator superalgebras and twisted modules, in “Mooshine, the Monster, and related topics” (C. Dong and G. Mason, Eds.), *Contemp. Math.* **193**, 203–236, Amer. Math. Soc., Providence, 1996.
- [Li3] H. Li, An analogue of the Hom functor and a generalized nuclear democracy theorem, *Duke Math. J.* **93** (1998), 73–114.
- [Ma] A. Matsuo, Norton’s trace formulae for the Griess algebra of a vertex operator algebra with larger symmetry, *Comm. Math. Phys.* **224** (2001), 565–591.
- [M1] M. Miyamoto, Griess algebras and conformal vectors in vertex operator algebras, *J. Algebra* **179** (1996), 528–548.
- [M2] M. Miyamoto, Binary codes and vertex operator (super)algebras, *J. Algebra* **181** (1996), 207–222.
- [M3] M. Miyamoto, Representation theory of code vertex operator algebras, *J. Algebra* **201** (1998), 115–150.
- [M4] M. Miyamoto, A Hamming code vertex operator algebra and construction of vertex operator algebras, *J. Algebra* **215** (1999), 509–530.
- [M5] M. Miyamoto, A new construction of the moonshine vertex operator algebra over the real number field, to appear in *Ann. of Math.*
- [M6] M. Miyamoto, Automorphism groups of \mathbb{Z}_2 -orbifold VOAs, preprint.
- [Sh] H. Shimakura, Automorphism group of the vertex operator algebra V_L^+ for an even lattice L without roots, math.QA/0311141.
- [SY] S. Sakuma and H. Yamauchi, Vertex operator algebra with two Miyamoto involutions generating S_3 , *J. Algebra* **267** (2003), 272–297.
- [Tu] M. Tuite, On the relationship between Monstrous Moonshine and the uniqueness of the Moonshine module, *Comm. Math. Phys.* **166** (1995), 495–532.
- [W] W. Wang, Rationality of Virasoro vertex operator algebras, *Internat. Math. Res. Notices* **71** (1993), 197–211.
- [Y1] H. Yamauchi, Module category of simple current extensions of vertex operator algebras, to appear in *J. Pure Appl. Algebra*.
- [Y2] H. Yamauchi, A theory of simple current extensions of vertex operator algebras and applications to the moonshine vertex operator algebra, to appear in Ph.D. thesis, University of Tsukuba, 2004.
- [Z] Y. Zhu, Modular invariance of characters of vertex operator algebras, *J. Amer. Math. Soc.* **9** (1996), 237–302.